

PROTEIN FAMILY REVIEW

Cyclin-dependent kinases

Marcos Malumbres

Summary

Cyclin-dependent kinases (CDKs) are protein kinases characterized by needing a separate subunit - a cyclin - that provides domains essential for enzymatic activity. CDKs play important roles in the control of cell division and modulate transcription in response to several extra- and intracellular cues. The evolutionary expansion of the CDK family in mammals led to the division of CDKs into three cell-cycle-related subfamilies (Cdk1, Cdk4 and Cdk5) and five transcriptional subfamilies (Cdk7, Cdk8, Cdk9, Cdk11 and Cdk20). Unlike the prototypical Cdc28 kinase of budding yeast, most of these CDKs bind one or a few cyclins, consistent with functional specialization during evolution. This review summarizes how, although CDKs are traditionally separated into cell-cycle or transcriptional CDKs, these activities are frequently combined in many family members. Not surprisingly, deregulation of this family of proteins is a hallmark of several diseases, including cancer, and drug-targeted inhibition of specific members has generated very encouraging results in clinical trials.

Gene organization and evolutionary history

Cyclin-dependent kinases (CDKs) are serine/threonine kinases whose activity depends on a regulatory subunit - a cyclin. Based on the sequence of the kinase domain, CDKs belong to the CMGC group of kinases (named for the initials of some members), along with mitogen-activated protein kinases (MAPKs), glycogen synthase kinase-3 beta (Gsk3 β), members of the dual-specificity tyrosine-regulated kinase (DYRK) family and CDK-like kinases [1]. In related kinases such as MAPKs, substrate specificity is conferred by docking sites separated from the catalytic site, whereas CDKs are characterized by dependency on separate protein subunits that provide additional sequences required for enzymatic activity. To aid nomenclature and analysis of CDKs, proteins belonging

to this family have been recently renamed as Cdk1 through to Cdk20 [2].

CDKs were first discovered by genetic and biochemical studies in model organisms such as yeasts and frogs (reviewed in [3]). This work established the importance of CDKs in promoting transitions through the cell cycle. In addition, these studies showed that the catalytic subunit, the CDK, must associate with a regulatory subunit, the cyclin, whose protein levels are subject to regulation during the cell cycle (this oscillation lent these regulators their cyclin name). Since these pioneer studies conducted in the 1980s, the importance of CDKs acting as a major eukaryotic protein kinase family involved in the integration of extracellular and intracellular signals to modulate gene transcription and cell division has been clearly established [3-6].

Despite their function in eukaryotic cell division and transcription, CDKs have undergone an extraordinary degree of evolutionary divergence and specialization. Six different CDKs are present in budding yeast (Figure 1). These CDKs can be grouped as, first, CDKs that bind multiple cyclins and can regulate the cell cycle and, second, CDKs that are activated by a single cyclin and are involved in the regulation of transcription. In the budding yeast *Saccharomyces cerevisiae*, the first group contains Cdc28 and Pho85, each binding nine or ten different cyclins, respectively. This promiscuity forms the basis for their dynamic regulation and their ability to phosphorylate multiple substrates, thus regulating the cell-division cycle in response to different cellular cues. The second group comprises four CDKs - Kin28, Srb10, Bur1 and Ctk1 - each activated by a single specific cyclin (Figure 1). These cyclins are usually not regulated in a cell-cycle-dependent manner, and the members of this second group of CDKs are involved in the control of gene transcription.

The number of CDKs increased during evolution and was marked by a greater expansion of the cell-cycle-related group. Fungi contain 6 to 8 CDKs and 9 to 15 cyclins, whereas flies and echinodermata contain 11 CDKs and 14 cyclins, and human cells have 20 CDKs and 29 cyclins (Box 1) [7]. Evolutionary studies suggest

Correspondence: malumbres@cniio.es
Cell Division and Cancer Group, Spanish National Cancer Research Centre (CNIO), Melchor Fernández Almagro 3, Madrid E-28029, Spain




Figure 1 Comparison of yeast and mammalian CDKs. Cells of the budding yeast *Saccharomyces cerevisiae* contain two cell-cycle-related CDKs that are activated by multiple cyclins - Cdc28 and Pho85. Cdk1 is the mammalian ortholog of Cdc28, whereas Cdk5 is considered to be the Pho85 ortholog. The Cdk4/Cdk6 subfamily is not present in yeast. Kin28, Srb10, Bur1 and Ctk1 are the yeast orthologs of Cdk7, Cdk8, Cdk9 and Cdk12, respectively. The Cdk20 and Cdk11/Cdk10 subfamilies are not represented in yeast. Also indicated is the cyclin partner for the mammalian CDKs. CDK, cyclin-dependent kinase.

that CDKs fall into eight subfamilies represented by Cdk1, Cdk4 and Cdk5 (from the yeast cell-cycle-related CDKs), and Cdk7, Cdk8, Cdk9, Cdk11 and Cdk20 (functioning as transcriptional CDKs) [7,8] (Figure 2). Like its yeast ortholog, Cdk1 is the only CDK essential for the cell cycle in mammals [9], whereas both Cdk2 and Cdk3 are dispensable [3,10]. Although Pho85 is not essential in yeast, this kinase is required for viability in some

stress conditions, such as growth after starvation. Pho85 displays multiple cell-cycle-related functions as well as regulation of gene expression, metabolism, morphogenesis, cell polarity and aging; it functions as an integrator of signals such as nutrient availability, DNA damage or other types of stress [11]. Sequencing and functional studies suggest that the mammalian homolog of Pho85 is Cdk5, although these kinases cluster with multiple mammalian kinases of the Cdk5 subfamily, namely Cdk14 to Cdk18. Pho85 can interact with up to 10 cyclins of the Pcl1/Pcl2 or Pho80 groups, whereas mammalian Cdk5 is activated by non-cyclin proteins, including Cdk5R1 (p35) and Cdk5R2 (p39). Interestingly, other members of the Cdk5 subfamily, such as Cdk14 or Cdk16, are activated by cyclin Y, which is a cyclin closely related to yeast Pcl1/Pcl2 proteins [12,13]. The Cdk4 subfamily is unique as it is only found in eumetazoans, and the members of this family diverge equally from the Cdk1 or Cdk5 subfamilies (Figures 1 and 2) [7]. Other cell-cycle-related subfamilies, such as the Cdk1-related B-type CDKs, are plant specific and are not found in animals or fungi [14].

Transcriptional CDKs are more conserved, both in sequence and function (Figure 1). Yeast Kin28 and human Cdk7 are subunits of transcription factor TFIIH, which is involved in transcription initiation by phosphorylating the Ser5 residue of the RNA polymerase II (RNAPII) C-terminal domain (CTD) at gene promoters. Cdk7 is also able to phosphorylate and activate other CDKs, thus acting as a CDK-activating kinase (CAK; Box 2). Kin28 does not have this activity, which is mediated in yeast by a different kinase unrelated to CDKs, Cak1 [8]. The yeast protein Srb10 is orthologous to human Cdk8 and Cdk19 and is the enzymatic component of the Mediator

Box 1. The cyclin family

Cyclins are a large family of approximately 30 proteins varying in mass from 35 to 90 kDa. These proteins are structurally defined by the presence of the so-called cyclin box, a domain of approximately 100 amino acid residues that forms a stack of five α -helices. Many cyclins have two cyclin boxes, one amino-terminal box for binding to CDKs, and a carboxy-terminal box that is usually required for the proper folding of the cyclin molecule. The cyclin box is also present in other molecules such as the retinoblastoma protein (Rb), the transcription factor TFIIIB and Cables (CDK5 and ABL1 enzyme substrate 1), which are unlikely to function as CDK activators. In general, cyclins show less sequence similarity than the CDKs. The cyclin family contains approximately 29 protein in humans, clustered in 16 subfamilies and three major groups: group I (cyclin B group: A-, B-, D-, E-, F-, G, J, I and O); group II (cyclin Y group - a partner of the Cdk5 subfamily); and group III (cyclin C group: C-, H-, K-, L- and T - major partners of transcriptional CDKs) [7,66]. Cyclin D and cyclin E clades (partners of Cdk1 and Cdk4 subfamilies) have undergone lineage-specific expansion and specialization in metazoa and plants [7].




Figure 2 Evolutionary relationships among the mammalian CDK subfamilies. The name of the different CDK subfamilies functioning in the cell cycle (orange) or transcription (green) is shown in boldface, and the domain structure of the individual proteins is depicted. The conserved protein kinase domain (red) and some additional domains (see key) are indicated for each CDK. Human cells contain two separate genes, *Cdk11A* and *Cdk11B*, each of them encoding a long isoform, Cdk11^{p110}, and a shorter protein, Cdk11^{p58}, generated by an internal ribosome binding site. The phylogenetic tree is based on the comparison of the human kinase domains [1]. CDK, cyclin-dependent kinase.

complex involved in the regulation of RNAPII during transcription [15]. Cdk9 is the yeast Bur1 ortholog, whereas the function of yeast Ctk1 in the phosphorylation of the RNAPII CTD is performed by Cdk12 in *Drosophila* and in human cells [16]. The evolutionary

relationship of the Cdk11 and Cdk20 subfamilies to the yeast CDKs is not clear, although these proteins are well conserved [7]. Unlike cyclins for cell-cycle-related kinases, the cyclin subunits of transcriptional CDKs do not show significant oscillations in protein levels during the cell cycle, and these transcriptional CDKs are therefore regulated by protein-protein interactions or other mechanisms. Transcription-related kinases possibly originated after cell-cycle-related CDKs and became more diverse as the complexity of transcription increased [17].

Box 2. The CDK-activating kinase complex

The CAK complex (comprising Cdk7, cyclin H and Mat1) phosphorylates the T-loop of all CDKs tested, thus participating in their activation. Furthermore, this complex can be part of the transcription factor phosphorylating the CTD of RNAPII as well as multiple nuclear receptors such as retinoic acid or thyroid receptors, the estrogen receptor α or the vitamin D receptor co-activator Ets1 [33]. The CAK complex can also be found associated with an additional subunit of TFIIH - the DNA-dependent helicase Xpd - forming a complex known as CAK-XPD. This complex plays a role in the coordination and progression of mitosis, likely as a consequence of the redistribution of CAK within different cell compartments during the late nuclear-division steps [67].

Characteristic structural features

Like other CMGC kinases, CDKs are proline-directed serine/threonine-protein kinases with some preference for the S/T-P-X-K/R sequence as a consequence of the presence of a hydrophobic pocket near the catalytic site that accommodates the proline (position +1). However, the requirement for the basic residue in the +3 position is not maintained in Cdk4 or transcriptional CDKs, which display a less-stringent S/T-P-X consensus. Some other family members such as Cdk7 or Cdk9 are not necessarily proline directed and can also phosphorylate residues in the absence of the +1 proline [18].

The CDKs range in size from approximately 250 amino acid residues, just encompassing the catalytic serine/threonine kinase domain, to proteins of more than 1,500 residues, with amino- and/or carboxy-terminal extensions of variable lengths (Figure 2). Like all kinases, CDKs have a two-lobed structure. The amino-terminal lobe contains beta-sheets, whereas the carboxy-terminal lobe is rich in α -helices, and the active site is sandwiched in-between. The N-lobe contains a glycine-rich inhibitory element (G-loop) and a unique major helix - the C-helix (containing the PSTAIRE sequence in Cdk1). The C-lobe contains the activation segment, which spans from the DFG motif (D145 in Cdk2; EMBL:AK291941) to the APE motif (E172 in Cdk2) and includes the phosphorylation-sensitive (T160 in Cdk2) residue in the so-called T-loop (Figure 3). In the cyclin-free monomeric form the CDK catalytic cleft is closed by the T-loop, preventing enzymatic activity. In addition, the activation segment in the C-lobe - a platform for binding of the phospho-acceptor Ser/Thr region of substrates - is partially disordered.

Cyclin-dependent kinase activation

Upon binding of the cyclin to Cdk2, the CDK C-helix packs against one specific helix in the cyclin partner through a surface characterized by extensive hydrophobic interactions. Association of cyclins to the C-helix promotes a rotation in the axis of this segment, generating new interactions that are part of the active ATP-binding site. In addition, cyclins take the C-lobe activation segment out of the catalytic site so that the threonine becomes accessible for activating

phosphorylation by CAK (Figure 3). This phosphothreonine acts as a rigidifying hub, stabilizing the activated form of the kinase heterodimer [18,19]. The extent of the CDK-cyclin interface varies in the structure of Cdk4, Cdk9 or yeast Pho85 [18,20,21]. For instance, Cdk2 and cyclin A contact each other at both the N- and C-lobes, whereas the contacts between Cdk4 and cyclin D are limited to the N-lobe, and, unlike Cdk2, the cyclin does not impose an active conformation on the kinase as the Cdk4 ATP-binding site is still inaccessible to its substrates, even in the presence of the cyclin [20,21]. How Cdk4 becomes active is not completely clear, although the binding of the substrate is thought to induce the activation segment to open and to fit to the phospho-acceptor site. Some CDKs, such as Cdk5 or its yeast ortholog Pho85, do not require phosphorylation in the activation segment for activity, and these kinase can adopt the correct conformation through other interactions [18].

In addition to the consensus kinase domain, a few CDKs contain additional domains with functional relevance. Cdk16, Cdk17 and Cdk18 (containing a PCTAIRE sequence in the C-helix) are characterized by a conserved catalytic domain flanked by amino- and carboxy-terminal extensions involved in cyclin binding. Phosphorylation of the Cdk16 amino-terminal domain blocks binding to cyclin Y, providing a novel mechanism for regulation of these complexes [22]. In Cdk12 and Cdk13 (characterized by a PITAIR motif), the kinase domain is localized in the center, and additional Arg/Ser-rich motifs in the amino terminus serve as docking sites for the assembly of splicing factors and regulators of splicing (Figure 2). These two kinases also contain




Figure 3 A three-dimensional view of CDK structure and activation. In monomeric Cdk2 (left; [PDB: 1HCL]), the major C-helix (N-lobe) and the activation domain are close, ensuring that the catalytic pocket is inaccessible. Upon binding of cyclin A (right; [PDB: 1JST]), the C-helix and the activation domain are pulled apart - a configuration that is further fixed by phosphorylation of residue T160, making the catalytic pocket accessible for enzymatic activity. The position of the inhibitory Thr14 (T14) and Tyr15 (Y15) residues in the G-loop is also shown. Color code: CDK subunit, orange; cyclin subunit, green; purple indicates specific named protein domains. CDK, cyclin-dependent kinase.

proline-rich motifs, mostly concentrated in their carboxy-terminal region, that serve as binding sites for Src-homology 3 (SH3), WW or profilin-domain-containing proteins [16].

Cyclin-dependent kinase inhibition

The glycine-rich region (G-loop) in the N-lobe is an additional regulatory region as it contains residues (Thr14 and Tyr15 in Cdk2; Figure 3) whose phosphorylation inhibits kinase activity. Phosphorylation of Thr14 and/or Tyr15 residues by Wee1 and Myt1 kinases inhibits several family members, preventing cell-cycle progression, for instance, in response to DNA damage. Elimination of these phosphates by phosphatases of the Cdc25 family is then required for activation of CDKs and cell-cycle progression [3,23]. Inhibitory phosphorylation at Thr14 and Tyr15 does not result in major changes in the CDK structure, but does inhibit the CDK activity by reducing the affinity of the CDK for its substrates. However, phosphorylation at Tyr15 seems to be activating in the case of Cdk5, perhaps by improving substrate recognition [18]. These residues are not present in Cdk7, in agreement with

the general belief that this kinase is constitutively active and regulated at different levels.

Cell-cycle-related CDKs can also be negatively regulated by binding to small proteins of the INK4 or Cip/Kip families of inhibitors [19,24]. INK4 proteins (p16^{INK4a}, p15^{INK4b}, p18^{INK4c} and p19^{INK4d}) are specific for the Cdk4 subfamily and interact with the monomeric CDKs. They function by distorting the cyclin interface and the ATP-binding pocket, thus preventing activation of Cdk4 and Cdk6 by D-type cyclins or by CAK [24]. Members of the Cip/Kip family of inhibitors (p21^{Cip1}, p27^{Kip1} and p57^{Kip2}) contact both the CDK and cyclin subunits and are able to inhibit CDK-cyclin heterodimers, giving additional levels of regulation once these complexes have already formed [19].

Localization and function

Cdk1 and Cdk4 subfamilies

The general picture in mammalian cells is that Cdk4 and Cdk6, upon transcriptional induction of D-type cyclins in response to several mitogenic stimuli, promote entry into the cell cycle (Figure 4) [25]. These kinases




Figure 4 An overview of CDK functions in the cell. Each CDK (in orange boxes) is shown in a complex with its major partner (green) - for clarity, only a few substrates are depicted. Most CDKs function in the nucleus (orange background), whereas a few family members are attached to the cell membrane or display cytoplasmic activities (blue background). Classical cell cycle CDKs - Cdk4, Cdk6, Cdk2 and Cdk1 - regulate the transitions through the different phases of the cell-division cycle. These activities are at least partially mediated by the control of multiple transcription factors (TFs) or regulatory elements such as the retinoblastoma protein (Rb). Cdk10 and Cdk11 also control transcription by phosphorylating TFs, hormone receptors and associated regulators (HRs), or splicing factors (SPFs). Cdk7, Cdk9 and Cdk12 directly phosphorylate the C-terminal domain (CTD) of RNA polymerase II (RNAPII), thus modulating the different phases of generation of transcripts. The Mediator complex is specifically regulated by Cdk8 or the highly related Cdk19. Cdk7 functions as a CDK-activating kinase (CAK) by directly phosphorylating several of the CDKs mentioned above. Cdk5 displays many functions in the cell, but it is better known for its function in the control of neuron-specific proteins such as Tau. The members of the Cdk14 subfamily, such as Cdk14 itself or Cdk16, are activated at the membrane by cyclin Y and also participate in many different pathways, such as Wnt-dependent signaling or signal transduction in the primary cilium. It is important to note that, for clarity, many interactions between CDKs and other partners, substrates or cellular processes are not shown - for instance, Cdk1 can bind to other cyclins and can also phosphorylate more than 100 substrates during mitotic entry that are not indicated here. CAK, CDK-activating kinase; CDK, cyclin-dependent kinase; CTD, C-terminal domain; Rb, retinoblastoma protein; RNAPII, RNA polymerase II; SPF, splicing factor; TF, transcription factor.

phosphorylate and inactivate the retinoblastoma protein (Rb), an adaptor protein that assembles different protein and protein-DNA complexes that repress transcription in response to a wide range of control mechanisms [25]. In human cells, Rb contains 13 conserved sites that are phosphorylated by CDKs in proliferating cells. Complexes between cyclin D and Cdk4 or Cdk6 phosphorylate residues Ser807 and Ser811, priming Rb for further phosphorylation by these or other CDKs at other sites [26]. CDK-dependent inactivation of Rb (or its relatives p107 and p130) results in de-repression of multiple genes encoding proteins required for DNA synthesis (S phase) or mitosis [25]. The activity of Cdk2 might also contribute to this process, although this kinase could have additional functions in DNA replication or DNA repair. Once cells have duplicated their DNA, Cdk1 becomes activated by A- and B-type cyclins, promoting cellular processes such as centrosome maturation and separation, chromosome condensation and mitotic entry after nuclear envelope breakdown [3]. This simplified view is obscured owing to multiple non-consensus interactions between CDKs and cyclins and compensatory roles [6]. For instance, when Cdk4 and Cdk6 are absent, Cdk2 can bind to D-type cyclins [27]. Cdk1 can also bind to cyclin E or cyclin D in the absence of Cdk2 or Cdk4, respectively [9], suggesting a scenario reminiscent of the yeast cell cycle in which Cdc28 is sufficient to induce all cell-cycle transitions by interacting with different cyclins [6].

Cdk5 subfamily and cyclin-Y-related kinases

Despite its similarity to other cell-cycle-related Cdk5 is the prototype of what are termed atypical CDKs. This kinase is activated by the non-cyclin proteins Cdk5R1 (p35) or Cdk5R2 (p39), and phosphorylation in the T-loop is not required for its activation [28,29]. Although Cdk5 is expressed in multiple cell types, its activity is thought to be more restricted owing to the expression of its activators p35 and p39 in terminally differentiated cells such as neurons [28]. However, in addition to its crucial functions in neuronal biology, Cdk5 plays multiple roles in gene expression, differentiation, angiogenesis and senescence, among others [5,28,29].

Interestingly, the Cdk5 activators carry an amino-terminal myristoylation motif that is required for their membrane targeting (Figure 4). Until recently, Cdk5 was thought to be the only membrane-associated Cdk, but recent data suggest that the CDKs Cdk14 to Cdk18 (PFTAIRE and PCTAIRE kinases) display similar activities upon binding to cyclin Y. Like Cdk5, Cdk16 requires no T-loop phosphorylation, suggesting that cyclin Y, like p35, tightly interacts with the activation loop, alleviating the need for an activating phosphorylation [13]. Cyclin Y is also N-myristoylated, and cyclin-Y-dependent recruitment

and activation of Cdk14 at the plasma membrane results in phosphorylation of the Wnt co-receptor Lrp5/Lrp6 (Figure 4). Cdk16 also binds to cyclin Y, and these complexes phosphorylate several proteins, including N-ethylmaleimide-sensitive factor (NSF) for the control of exocytosis [30], and are essential for spermatogenesis [22]. The partner CDKs of cyclin Y display overlapping roles as knockdown of individual CDKs in *Xenopus* embryos failed to produce a phenotype, whereas depletion of cyclin Y and its highly related homolog cyclin-Y-like resulted in a Wnt loss-of-function phenotype [31]. In addition to the relevance of the Wnt pathway in the control of transcription, β -catenin and other Wnt regulators localize to centrosomes and/or kinetochores and regulate the formation and orientation of the mitotic spindle and the process of chromosome segregation [31]. In fact, cyclin Y reaches maximum levels at G2-M phase of the cell cycle and is degraded in a ubiquitin-dependent manner, similarly to mitotic cyclins, suggesting a crucial role for the cyclin-Y-Wnt pathway during cell division [12]. It is interesting to note that CDKs and cyclins of this subfamily, such as Cdk17 or cyclin Y, are highly conserved, at levels similar to Cdk1 or cyclin B [13]. In most cases, the cellular relevance of many Cdk5-subfamily members remains to be established.

Control of RNA polymerase II by transcriptional cyclin-dependent kinases

One of the most important activities of CDKs is reversible phosphorylation of the CTD of the largest subunit (Rpb1) of RNAPII (Figure 4). The CTD consists of multiple repeats of an evolutionarily conserved heptapeptide possessing the consensus sequence Tyr-Ser-Pro-Thr-Ser-Pro-Ser, with the number of repeats varying among different organisms, ranging from 26 repeats in yeast to 52 in mammals. The CTD is the target of multiple post-translational modifications, including phosphorylation, generating a complex regulatory code known as the CTD code. The CTD regulates the cycling of RNAPII between a hypophosphorylated form, able to enter the preinitiation complex, and a hyperphosphorylated form capable of processive elongation of the transcript [32]. Multiple CDKs can phosphorylate the CTD, including cell-cycle-related kinases Cdk1 or Cdk2 and most transcriptional CDKs of the Cdk7, Cdk8 and Cdk9 subfamilies (Figure 4). Cdk7 is a member of the ten-subunit general transcription factor TFIIB^b that phosphorylates Ser5 and Ser7 of the heptad during initiation and promoter clearance [33,34]. Cdk7 also phosphorylates and activates Cdk9, thus promoting downstream events [34]. To release the paused RNAPII and allow productive elongation, Ser2 of the heptad is then phosphorylated, a process in which both Cdk9 and Cdk12 have been implicated. Cdk9 binds to T-type cyclins (T1 and T2) as a

subunit of the positive transcription elongation factor b (P-TEFb) that stimulates elongation. Cdk9 is the ortholog of Bur1, which contributes to phosphorylation of the Ser2 mark at the 5' ends of genes [16,35]. Although Cdk9 was thought to be the major Ser2 kinase required for efficient elongation, recent data suggest that this requirement is mediated by a second substrate of Cdk9, the elongation factor subunit Spt5, whose Cdk9-dependent phosphorylation relieves the early pausing step [35]. Recent studies in *Drosophila* and human cells suggest that Cdk12, in complex with cyclin K, is the yeast Ctk1 ortholog responsible for most of the Ser2 phosphorylation at the CTD and especially the phosphorylation at promoter-distal regions [36,37]. Depletion of Cdk12 resulted in defective Ser2 phosphorylation at a subset of genes - mostly long and complex ones - but not a change in the rate of global transcription. Cdk12 is specifically required for the transcription of genes involved in the response to DNA damage, establishing a new link between the transcriptional machinery and cell-cycle regulation [37]. Cdk1 can also phosphorylate the CTD, and this activity is thought to inhibit transcription, although its physiological relevance has not been established. Transcript termination results in dephosphorylation of RNAPII, making it ready for another round of re-initiation. Although the control of dephosphorylation is not well understood, several CDK-counteracting phosphatases such as Cdc14 are likely to be involved [38,39].

Cdk8 and its closely related family member Cdk19 associate with C-type cyclins as part of the multi-subunit Mediator complex (Figure 4) [15]. This complex functions as a bridge linking gene-specific activators to the general RNAPII transcription machinery at the promoter, thus influencing nearly all stages of transcription and coordinating these events with changes in chromatin organization. Cdk8 (or Cdk19), along with cyclin C, Med12 and Med13, form the so-called Cdk8 module characteristic of the free Mediator form, devoid of RNAPII. The Cdk8 module responds to several intracellular signaling pathways, and it is commonly associated with repression of transcription, although it can also activate transcription [15]. Cdk8 has multiple targets and phosphorylates several transcription factors, affecting their stability and activity. Recent evidence suggests various roles in gene activation in the p53 network, the Wnt- β -catenin pathway, the serum-response network and other pathways governed by Smads or the thyroid hormone receptor [40]. Cdk8 also modulates Cdk7 activity by phosphorylating cyclin H, thus impeding Cdk7 activity and inhibiting initiation of transcription [33]. Finally, Cdk19 associates with similar Mediator complexes, although these complexes are likely to possess a specificity that is yet to be established [41].

Cdk11 and Cdk20 subfamilies

Cdk11 proteins are the products of two highly related genes in mammals (*CDK11A* and *CDK11B*) encoding Cdk11A^{P110} and Cdk11B^{P110} [2], as well as two smaller alternative proteins, Cdk11A^{P58} and Cdk11B^{P58}, resulting from translation from an internal ribosome-binding site generated during G2-M phase. Cdk11 binds to L-type cyclins and participates in the coordination between transcription and RNA processing, particularly alternative splicing [42]. In budding yeast, Cdk11 has been shown to be a crucial factor for the interaction of the Cdk8 module with the Mediator complex through phosphorylation of conserved residues of the Med27 and Med4 Mediator subunits (Figure 4) [43]. Cdk11 also participates in many other pathways, such as hormone receptor signaling or autophagy [44-46]. The short isoform of Cdk11, Cdk11^{P58}, is specifically expressed at G2-M, and its kinase activity is required for duplication of the centrioles, spindle dynamics and sister chromatid cohesion at centromeres during mitosis [47-49]. Lack of Cdk11 results in mitotic defects in mouse embryos, highlighting the crucial role of this 'transcriptional' kinase in the cell cycle [3].

Cdk10 is activated by cyclin M, a cyclin mutated in STAR syndrome, a developmental abnormality characterized by toe syndactyly, telecanthus and anogenital and renal malformations [50]. Cdk10-cyclin-M phosphorylates Ets2, promoting its degradation by the proteasome [50]. STAR-associated mutations in the gene encoding cyclin M impair binding of cyclin M to Cdk10, resulting in increased Ets2-dependent transcription of c-Raf and over-activation of the MAPK pathway. In the insect *Helicoverpa armigera*, Cdk10 modulates gene transcription by steroid hormones by promoting the interaction between heat-shock proteins and the ecdysone receptor EcRB1 [51].

Finally, Cdk20 (also known as cell cycle-related kinase (CCRK)) can interact with cyclin H and originally was proposed to have CAK activity for Cdk2, suggesting a close relationship with Cdk7. However, its role as a CAK is controversial [52], and additional data suggest that it functions as an activating kinase for MAK-related kinase/intestinal cell kinase (ICK) [53]. Expression of Cdk20 activates β -catenin-TCF signaling to stimulate cell-cycle progression [54], whereas its inhibition results in accumulation of ICK at the ciliary tips and prevents cell-cycle entry [55] (Figure 4).

Frontiers

It is abundantly clear that the CDK family is central to multiple signaling pathways controlling transcription and cell-cycle progression. CDKs probably originated as a system to modulate cell-cycle-promoting activity in response to various cellular scenarios. Over the course of

evolution, both CDK and cyclin gene families have independently undergone a significant number of functional specializations [7]. Many of the interactions between specific mammalian CDKs and cyclins have been reported *in vitro*. However, the biochemical promiscuity in CDK-cyclin interactions makes it difficult to evaluate properly the *in vivo* physiological relevance of specific CDK-cyclin complexes. For instance, Cdk1 is thought to be activated mainly by A- and B-type cyclins but can also bind to, and be activated by, D- or E-type cyclins in the absence of Cdk4/Cdk6 or Cdk2, respectively [9,27,56]. Cdk5 can also bind to D-type cyclins, although to what extent these complexes are active or relevant *in vivo* is not clear. The situation is even more complex for the lesser-known family members for which there are no current *in vivo* data [2].

Although the comparison of the yeast CDKs has promoted the convenient division between transcriptional and cell-cycle activities, the multiple interactions between these two activities in higher eukaryotes makes it difficult to maintain this simple classification. First, transcription and cell-cycle progression cannot be opposed as these processes function at different layers in cell biology. Arguably, transcription is a major regulatory pathway required for cell-cycle entry. Major cell-cycle-related kinases such as Cdk4 and Cdk6 mostly function by phosphorylating transcription regulators such as Rb or Smads [3,25], and the archetypal cell-cycle kinase Cdk1 also phosphorylates multiple transcription factors and epigenetic modulators (Figure 4) [5]. By contrast, major 'transcriptional' CDKs such as Cdk7 or Cdk11 directly control cell-cycle progression, in some instances independently of transcription. Finally, a single CDK can have separate cell-cycle-related and transcriptional activities. As an example, Cdk6 has recently been characterized as a chromatin factor (Figure 4) that regulates transcription factors involved in angiogenesis or the NF- κ B pathway [57,58], a process independent of the classical Cdk4/6-cyclin-D-Rb pathway involved in cell-cycle regulation.

As a consequence of their importance in multiple processes, CDKs are frequently mutated or deregulated in disease. A classic example is the almost universal deregulation of the CDK-cyclin-Rb pathway in cell-cycle entry during malignant transformation [25]. Underlining the significance of CDKs, inhibitors of Cdk4 and Cdk6 received in 2013 the Food and Drug Administration 'breakthrough therapy' designation for treatment of patients with breast cancer [59]. Other members of the CDK family can also be considered as interesting targets for therapeutics in cancer or other diseases. Cdk5 displays multiple roles in neurodegenerative diseases [28] and in other tissues with relevance to diabetes, cardiovascular disease or cancer [29]. Cdk8 exhibits copy-number gains in colon cancers, and recently it has been characterized as a coactivator of the

beta-catenin pathway in colon cancer cell proliferation [60,61]. Cdk10 is a major determinant of resistance to endocrine therapy for breast cancer [62], and inhibition of Cdk12 confers sensitivity to inhibitors of poly (ADP-ribose) polymerases PARP1 and PARP2 [63]. Cdk14 confers motility advantages and metastatic potential in hepatocellular carcinoma motility and metastasis [64,65]. Finally, as indicated above, cyclin Y kinases regulate the Wnt pathway [31], providing new therapeutic opportunities that are yet to be explored. Hence, it seems very likely that new targets within the CDK family will be explored in the near future for therapy of cancer or other diseases.

Abbreviations

CAK: CDK-activating kinase; CCRK: Cycle-related kinase; CDK: Cyclin-dependent kinase; CTD: C-terminal domain; DYRK: Dual-specificity tyrosine-regulated kinase; ICK: Intestinal cell kinase; MAPK: Mitogen-activated protein kinase; NSF: N-ethylmaleimide-sensitive factor; PARP: Poly (ADP-ribose) polymerase; P-TEFb: Positive transcription elongation factor b; Rb: Retinoblastoma protein; RNAPII: RNA polymerase II.

Competing interests

The author declares that he has no competing interests.

Acknowledgements

Work in the laboratory of MM is funded by grants from the Spanish Government (MINECO; SAF2012-38215), the OncoCycle Programme (S2010/BMD-2470) from the Comunidad de Madrid, and the European Union Seventh Framework Programme (MitoSys project; HEALTH-F5-2010-241548).

Published: 30 June 2014

References

- Manning G, Whyte DB, Martinez R, Hunter T, Sudarsanam S: **The protein kinase complement of the human genome.** *Science* 2002, **298**:1912–1934.
- Malumbres M, Harlow E, Hunt T, Hunter T, Lahti JM, Manning G, Morgan DO, Tsai LH, Wolgemuth DJ: **Cyclin-dependent kinases: a family portrait.** *Nat Cell Biol* 2009, **11**:1275–1276.
- Malumbres M, Barbacid M: **Mammalian cyclin-dependent kinases.** *Trends Biochem Sci* 2005, **30**:630–641.
- Morgan DO: **Cyclin-dependent kinases: engines, clocks, and microprocessors.** *Annu Rev Cell Dev Biol* 1997, **13**:261–291.
- Lim S, Kaldis P: **Cdks, cyclins and CKIs: roles beyond cell cycle regulation.** *Development* 2013, **140**:3079–3093.
- Malumbres M, Barbacid M: **Cell cycle, CDKs and cancer: a changing paradigm.** *Nat Rev Cancer* 2009, **9**:153–166.
- Cao L, Chen F, Yang X, Xu W, Xie J, Yu L: **Phylogenetic analysis of CDK and cyclin proteins in premetazoan lineages.** *BMC Evol Biol* 2014, **14**:10.
- Liu J, Kipreos ET: **Evolution of cyclin-dependent kinases (CDKs) and CDK-activating kinases (CAKs): differential conservation of CAKs in yeast and metazoa.** *Mol Biol Evol* 2000, **17**:1061–1074.
- Santamaria D, Barriere C, Cerqueira A, Hunt T, Tardy C, Newton K, Caceres JF, Dubus P, Malumbres M, Barbacid M: **Cdk1 is sufficient to drive the mammalian cell cycle.** *Nature* 2007, **448**:811–815.
- Ortega S, Prieto I, Odajima J, Martin A, Dubus P, Sotillo R, Barbero JL, Malumbres M, Barbacid M: **Cyclin-dependent kinase 2 is essential for meiosis but not for mitotic cell division in mice.** *Nat Genet* 2003, **35**:25–31.
- Huang D, Friesen H, Andrews B: **Pho85, a multifunctional cyclin-dependent protein kinase in budding yeast.** *Mol Microbiol* 2007, **66**:303–314.
- Davidson G, Shen J, Huang YL, Su Y, Karaulanov E, Bartscherer K, Hassler C, Stanek P, Boutros M, Niehrs C: **Cell cycle control of wnt receptor activation.** *Dev Cell* 2009, **17**:788–799.
- Mikolcovic P, Rainer J, Geley S: **Orphan kinases turn eccentric: a new class of cyclin Y-activated, membrane-targeted CDKs.** *Cell Cycle* 2012, **11**:3758–3768.
- Harashima H, Dissmeyer N, Schnittger A: **Cell cycle control across the eukaryotic kingdom.** *Trends Cell Biol* 2013, **23**:345–356.

15. Carlsten JO, Zhu X, Gustafsson CM: **The multitasking Mediator complex.** *Trends Biochem Sci* 2013, **38**:531–537.
16. Kohoutek J, Blazek D: **Cyclin K goes with Cdk12 and Cdk13.** *Cell Div* 2012, **7**:12.
17. Guo Z, Stiller JW: **Comparative genomics of cyclin-dependent kinases suggest co-evolution of the RNAP II C-terminal domain and CTD-directed CDKs.** *BMC Genomics* 2004, **5**:69.
18. Echalié A, Endicott JA, Noble ME: **Recent developments in cyclin-dependent kinase biochemical and structural studies.** *Biochim Biophys Acta* 1804, **2010**:511–519.
19. Pavletich NP: **Mechanisms of cyclin-dependent kinase regulation: structures of Cdk5, their cyclin activators, and Cip and INK4 inhibitors.** *J Mol Biol* 1999, **287**:821–828.
20. Day PJ, Cleasby A, Tickle IJ, O'Reilly M, Coyle JE, Holding FP, McMenamin RL, Yon J, Chopra R, Lengauer C, Jhoti H: **Crystal structure of human CDK4 in complex with a D-type cyclin.** *Proc Natl Acad Sci U S A* 2009, **106**:4166–4170.
21. Takaki T, Echalié A, Brown NR, Hunt T, Endicott JA, Noble ME: **The structure of CDK4/cyclin D3 has implications for models of CDK activation.** *Proc Natl Acad Sci U S A* 2009, **106**:4171–4176.
22. Mikolcevic P, Sigl R, Rauch V, Hess MW, Pfaller K, Barisic M, Pelliniemi LJ, Boesl M, Geley S: **Cyclin-dependent kinase 16/PCTAIRE kinase 1 is activated by cyclin Y and is essential for spermatogenesis.** *Mol Cell Biol* 2012, **32**:868–879.
23. Boutros R, Lobjois V, Ducommun B: **CDC25 phosphatases in cancer cells: key players? Good targets?** *Nat Rev Cancer* 2007, **7**:495–507.
24. Jeffrey PD, Tong L, Pavletich NP: **Structural basis of inhibition of CDK-cyclin complexes by INK4 inhibitors.** *Genes Dev* 2000, **14**:3115–3125.
25. Malumbres M, Barbacid M: **To cycle or not to cycle: a critical decision in cancer.** *Nat Rev Cancer* 2001, **1**:222–231.
26. Rubin SM: **Deciphering the retinoblastoma protein phosphorylation code.** *Trends Biochem Sci* 2013, **38**:12–19.
27. Malumbres M, Sotillo R, Santamaria D, Galan J, Cerezo A, Ortega S, Dubus P, Barbacid M: **Mammalian cells cycle without the D-type cyclin-dependent kinases Cdk4 and Cdk6.** *Cell* 2004, **118**:493–504.
28. Cheung ZH, Ip NY: **Cdk5: a multifaceted kinase in neurodegenerative diseases.** *Trends Cell Biol* 2012, **22**:169–175.
29. Arif A: **Extraneuronal activities and regulatory mechanisms of the atypical cyclin-dependent kinase Cdk5.** *Biochem Pharmacol* 2012, **84**:985–993.
30. Liu Y, Cheng K, Gong K, Fu AK, Ip NY: **Pctaire1 phosphorylates N-ethylmaleimide-sensitive fusion protein: implications in the regulation of its hexamerization and exocytosis.** *J Biol Chem* 2006, **281**:9852–9858.
31. Davidson G, Niehrs C: **Emerging links between CDK cell cycle regulators and Wnt signaling.** *Trends Cell Biol* 2010, **20**:453–460.
32. Egloff S, Dienstbier M, Murphy S: **Updating the RNA polymerase CTD code: adding gene-specific layers.** *Trends Genet* 2012, **28**:333–341.
33. Egly JM, Coin F: **A history of TFIIH: two decades of molecular biology on a pivotal transcription/repair factor.** *DNA Repair (Amst)* 2011, **10**:714–721.
34. Larochelle S, Amat R, Glover-Cutter K, Sanso M, Zhang C, Allen JJ, Shokat KM, Bentley DL, Fisher RP: **Cyclin-dependent kinase control of the initiation-to-elongation switch of RNA polymerase II.** *Nat Struct Mol Biol* 2012, **19**:1108–1115.
35. Bres V, Yoh SM, Jones KA: **The multi-tasking P-TEFb complex.** *Curr Opin Cell Biol* 2008, **20**:334–340.
36. Bartkowiak B, Liu P, Phatnani HP, Fuda NJ, Cooper JJ, Price DH, Adelman K, Lis JT, Greenleaf AL: **CDK12 is a transcription elongation-associated CTD kinase, the metazoan ortholog of yeast Ctk1.** *Genes Dev* 2010, **24**:2303–2316.
37. Blazek D, Kohoutek J, Bartholomeeusen K, Johansen E, Hulinkova P, Luo Z, Cimermanic P, Ule J, Peterlin BM: **The Cyclin K/Cdk12 complex maintains genomic stability via regulation of expression of DNA damage response genes.** *Genes Dev* 2011, **25**:2158–2172.
38. Clemente-Blanco A, Sen N, Mayan-Santos M, Sacristan MP, Graham B, Jarmuz A, Giess A, Webb E, Game L, Eick D, Bueno A, Merckenschlager M, Aragon L: **Cdc14 phosphatase promotes segregation of telomeres through repression of RNA polymerase II transcription.** *Nat Cell Biol* 2011, **13**:1450–1456.
39. Guillaumot M, Manchado E, Chiesa M, Gomez-Lopez G, Pisano DG, Sacristan MP, Malumbres M: **Cdc14b regulates mammalian RNA polymerase II and represses cell cycle transcription.** *Sci Rep* 2011, **1**:189.
40. Galbraith MD, Donner AJ, Espinosa JM: **CDK8: a positive regulator of transcription.** *Transcription* 2010, **1**:4–12.
41. Sato S, Tomomori-Sato C, Parmely TJ, Florens L, Zybailov B, Swanson SK, Banks CA, Jin J, Cai Y, Washburn MP, Conaway JW, Conaway RC: **A set of consensus mammalian mediator subunits identified by multidimensional protein identification technology.** *Mol Cell* 2004, **14**:685–691.
42. Hu D, Mayeda A, Trembley JH, Lahti JM, Kidd VJ: **CDK11 complexes promote pre-mRNA splicing.** *J Biol Chem* 2003, **278**:8623–8629.
43. Drogat J, Migeot V, Mommaerts E, Mullier C, Dieu M, van Bakel H, Hermand D: **Cdk11-cyclinL controls the assembly of the RNA polymerase II mediator complex.** *Cell Rep* 2012, **2**:1068–1076.
44. Wilkinson S, Croft DR, O'Prey J, Meedendorp A, O'Prey M, Dufes C, Ryan KM: **The cyclin-dependent kinase PITSLRE/CDK11 is required for successful autophagy.** *Autophagy* 2011, **7**:1295–1301.
45. Wang Y, Zong H, Chi Y, Hong Y, Yang Y, Zou W, Yun X, Gu J: **Repression of estrogen receptor alpha by CDK11p58 through promoting its ubiquitin-proteasome degradation.** *J Biochem* 2009, **145**:331–343.
46. Chi Y, Hong Y, Zong H, Wang Y, Zou W, Yang J, Kong X, Yun X, Gu J: **CDK11p58 represses vitamin D receptor-mediated transcriptional activation through promoting its ubiquitin-proteasome degradation.** *Biochem Biophys Res Commun* 2009, **386**:493–498.
47. Petretti C, Savoian M, Montembault E, Glover DM, Prigent C, Giet R: **The PITSLRE/CDK11p58 protein kinase promotes centrosome maturation and bipolar spindle formation.** *EMBO Rep* 2006, **7**:418–424.
48. Hu D, Valentine M, Kidd VJ, Lahti JM: **CDK11(p58) is required for the maintenance of sister chromatid cohesion.** *J Cell Sci* 2007, **120**:2424–2434.
49. Yokoyama H, Gruss OJ, Rybina S, Caudron M, Schelder M, Wilm M, Mattaj JW, Karsenti E: **Cdk11 is a RanGTP-dependent microtubule stabilization factor that regulates spindle assembly rate.** *J Cell Biol* 2008, **180**:867–875.
50. Guen VJ, Gamble C, Flajolet M, Unger S, Thollet A, Ferandin Y, Superti-Furga A, Cohen PA, Meijer L, Colas P: **CDK10/cyclin M is a protein kinase that controls ETS2 degradation and is deficient in STAR syndrome.** *Proc Natl Acad Sci U S A* 2013, **110**:19525–19530.
51. Liu W, Cai MJ, Wang JX, Zhao XF: **In a non-genomic action, steroid hormone 20-hydroxycyclopentanone induces phosphorylation of cyclin-dependent kinase 10 to promote gene transcription.** *Endocrinology* 2014, **155**:1738–1750.
52. Wohlbold L, Larochelle S, Liao JC, Livshits G, Singer J, Shokat KM, Fisher RP: **The cyclin-dependent kinase (CDK) family member PNQALRE/CCRK supports cell proliferation but has no intrinsic CDK-activating kinase (CAK) activity.** *Cell Cycle* 2006, **5**:546–554.
53. Fu Z, Larson KA, Chitta RK, Parker SA, Turk BE, Lawrence MW, Kaldis P, Galaktionov K, Cohn SM, Shabanowitz J, Hunt DF, Sturgill TW: **Identification of yin-yang regulators and a phosphorylation consensus for male germ cell-associated kinase (MAK)-related kinase.** *Mol Cell Biol* 2006, **26**:8639–8654.
54. Feng H, Cheng AS, Tsang DP, Li MS, Go MY, Cheung YS, Zhao GJ, Ng SS, Lin MC, Yu J, Lai PB, To KF, Sung JJ: **Cell cycle-related kinase is a direct androgen receptor-regulated gene that drives beta-catenin/T cell factor-dependent hepatocarcinogenesis.** *J Clin Invest* 2011, **121**:3159–3175.
55. Yang Y, Roine N, Makela TP: **CCRK depletion inhibits glioblastoma cell proliferation in a cilium-dependent manner.** *EMBO Rep* 2013, **14**:741–747.
56. Aleem E, Kiyokawa H, Kaldis P: **Cdc2-cyclin E complexes regulate the G1/S phase transition.** *Nat Cell Biol* 2005, **7**:831–836.
57. Kollmann K, Heller G, Schneckenleithner C, Warsch W, Scheicher R, Ott RG, Schafer M, Fajmann S, Schleiderer M, Schiefer AI, Reichart U, Mayerhofer M, Hoeller C, Zochbauer-Muller S, Kerjaschki D, Bock C, Kenner L, Hoeffler G, Freissmuth M, Green AR, Moriggl R, Busslinger M, Malumbres M, Sexl V: **A kinase-independent function of CDK6 links the cell cycle to tumor angiogenesis.** *Cancer Cell* 2013, **24**:167–181.
58. Handschick K, Beuerlein K, Jurida L, Bartkuhn M, Muller H, Soelch J, Weber A, Dietrich-Breiholz O, Schneider H, Scharfe M, Jarek M, Stellzig J, Schmitz ML, Kracht M: **Cyclin-dependent kinase 6 is a chromatin-bound cofactor for NF-kappaB-dependent gene expression.** *Mol Cell* 2014, **53**:193–208.
59. **Breakthrough therapies.** <http://www.focr.org/breakthrough-therapies>.
60. Firestein R, Bass AJ, Kim SY, Dunn IF, Silver SJ, Guney I, Freed E, Ligon AH, Vena N, Ogino S, Chheda MG, Tamayo P, Finn S, Shrestha Y, Boehm JS, Jain S, Bojarski E, Mermel C, Barretina J, Chan JA, Baselga J, Taberner J, Root DE, Fuchs CS, Loda M, Shivdasani RA, Meyerson M, Hahn WC: **CDK8 is a colorectal cancer oncogene that regulates beta-catenin activity.** *Nature* 2008, **455**:547–551.
61. Morris EJ, Ji JY, Yang F, Di Stefano L, Herr A, Moon NS, Kwon EJ, Haigis KM, Naar AM, Dyson NJ: **E2F1 represses beta-catenin transcription and is antagonized by both pRB and CDK8.** *Nature* 2008, **455**:552–556.

62. Iorns E, Turner NC, Elliott R, Syed N, Garrone O, Gasco M, Tutt AN, Crook T, Lord CJ, Ashworth A: **Identification of CDK10 as an important determinant of resistance to endocrine therapy for breast cancer.** *Cancer Cell* 2008, **13**:91–104.
63. Bajrami I, Frankum JR, Konde A, Miller RE, Rehman FL, Brough R, Campbell J, Sims D, Rafiq R, Hooper S, Chen L, Kozarewa I, Assiotis I, Fenwick K, Natrajan R, Lord CJ, Ashworth A: **Genome-wide profiling of genetic synthetic lethality identifies CDK12 as a novel determinant of PARP1/2 inhibitor sensitivity.** *Cancer Res* 2014, **74**:287–297.
64. Leung WK, Ching AK, Chan AW, Poon TC, Mian H, Wong AS, To KF, Wong N: **A novel interplay between oncogenic PFTK1 protein kinase and tumor suppressor TAGLN2 in the control of liver cancer cell motility.** *Oncogene* 2011, **30**:4464–4475.
65. Huang J, Deng Q, Wang Q, Li KY, Dai JH, Li N, Zhu ZD, Zhou B, Liu XY, Liu RF, Fei QL, Chen H, Cai B, Xiao HS, Qin LX, Han ZG: **Exome sequencing of hepatitis B virus-associated hepatocellular carcinoma.** *Nat Genet* 2012, **44**:1117–1121.
66. Ma Z, Wu Y, Jin J, Yan J, Kuang S, Zhou M, Zhang Y, Guo AY: **Phylogenetic analysis reveals the evolution and diversification of cyclins in eukaryotes.** *Mol Phylogenet Evol* 2013, **66**:1002–1010.
67. Chen J, Larochelle S, Li X, Suter B: **Xpd/Ercc2 regulates CAK activity and mitotic progression.** *Nature* 2003, **424**:228–232.

doi:10.1186/gb4184

Cite this article as: Malumbres: Cyclin-dependent kinases. *Genome Biology* 2014 **15**:122.

The history and future of targeting cyclin-dependent kinases in cancer therapy

Uzma Asghar¹, Agnieszka K. Witkiewicz², Nicholas C. Turner³ and Erik S. Knudsen²

Abstract | Cancer represents a pathological manifestation of uncontrolled cell division; therefore, it has long been anticipated that our understanding of the basic principles of cell cycle control would result in effective cancer therapies. In particular, cyclin-dependent kinases (CDKs) that promote transition through the cell cycle were expected to be key therapeutic targets because many tumorigenic events ultimately drive proliferation by impinging on CDK4 or CDK6 complexes in the G1 phase of the cell cycle. Moreover, perturbations in chromosomal stability and aspects of S phase and G2/M control mediated by CDK2 and CDK1 are pivotal tumorigenic events. Translating this knowledge into successful clinical development of CDK inhibitors has historically been challenging, and numerous CDK inhibitors have demonstrated disappointing results in clinical trials. Here, we review the biology of CDKs, the rationale for therapeutically targeting discrete kinase complexes and historical clinical results of CDK inhibitors. We also discuss how CDK inhibitors with high selectivity (particularly for both CDK4 and CDK6), in combination with patient stratification, have resulted in more substantial clinical activity.

Fundamentally, the cell cycle process is conserved from unicellular eukaryotes to complex metazoans¹, and distinct phases of the cell cycle are responsive to physiological cues that dictate the appropriateness of cell division. Cyclin-dependent kinases (CDKs) are critical regulatory enzymes that drive all cell cycle transitions^{1–6}, and their activity is under stringent control to ensure successful cell division. In particular, all mitotic cell division requires that faithful DNA replication occurs in S phase and that the requisite machinery to divide chromosomes is in place during mitosis, leading to the production of daughter cells. In unicellular eukaryotes, cell cycle progression is predominantly controlled by the availability of nutrients to ensure the completion of successful duplication. Cell cycle progression in unicellular eukaryotes is also dependent on the absence of genetic damage that would preclude the viability of daughter cells. In multicellular organisms, more complex regulatory mechanisms that reflect cell–cell communication have evolved.

Many of the key concepts of CDK biology (FIG. 1) were discovered >20 years ago through the study of yeast and the synchronous cycles of division seen in embryo extracts; indeed, the findings from studies led to the award of a Nobel Prize for these researchers^{7,8}.

In particular, CDK1 emerged as a key determinant of mitotic progression, and CDK2 emerged as being more relevant for DNA replication in higher eukaryotes. In metazoans, much of the control over cell cycle entry is elicited at the level of CDK4 and CDK6, which are responsive to numerous growth regulatory signals. Subsequently, in addition to the CDKs that directly promote cell cycle progression (for example, CDK4, CDK6, CDK2 and CDK1), an additional family of CDKs that regulate transcription was identified, which include CDK7, CDK8 and CDK9 (REFS 3,9–11). CDKs with post-mitotic functions in specialized tissue settings, such as CDK5, were also identified. Owing to the central role of CDKs in the control of cell division, it is perhaps not surprising that all cancers exhibit some features that derange the normal controls over the cell cycle¹², and over the past 20 years, numerous drugs that target CDK activity have emerged and have been tested in the clinic. Here, we review the biology of CDKs and their suitability as therapeutic targets in cancer, the key mechanisms through which CDKs become deranged in cancer and the challenges that have, until recently, complicated attempts to bring CDK inhibitors through to successful clinical application.

¹Breakthrough Breast Cancer Research Centre, Chester Beatty Laboratories, Institute of Cancer Research, London, SW3 6JB, UK.

²Simmons Cancer Center and Department of Pathology, University of Texas Southwestern, Dallas, USA.

³Institute of Cancer Research and Royal Marsden NHS Foundation Trust Breast Cancer Unit, London, SW3 6JJ, UK.

Correspondence to E.S.K.
e-mail: erik.knudsen@utsouthwestern.edu
doi:10.1038/nrd4504




Figure 1 | Progression of the cell cycle driven by CDKs. Mitogenic signals stimulate cyclin-dependent kinase 4 (CDK4) and CDK6 and promote entry into the cell cycle, whereas antiproliferative checkpoints inhibit CDK4 and CDK6 activity or induce the expression of the CDK4 and CDK6 inhibitor p16^{INK4A}. Active CDK4 and CDK6 complexes initiate the phosphorylation (P) of key substrates, including the tumour suppressor retinoblastoma protein (RB), thereby unleashing a gene expression programme that is coordinated by the E2F family of transcription factors. In this context, CDK4 and CDK6 initiate transcription and stability of E-type and A-type cyclins (CycE and CycA, respectively) and the subsequent activation of CDK2 that contributes to the further phosphorylation of RB and the initiation of DNA replication. Further checkpoints can directly inhibit CDK2 activity or induce the CDK-interacting protein/kinase inhibitory protein (CIP/KIP) class of inhibitors (p21^{CIP1} and p27^{KIP1}) that bind to and inhibit CDK2–cyclin complexes. With the completion of DNA replication, CDK1–Cyc A and CDK1–Cyc B complexes form to phosphorylate targets in G2 phase. In the absence of DNA damage and following appropriate preparation for chromosomal segregation, the cellular default is to activate CDK1–CycB complexes and progress into mitosis. However, there are potent checkpoints that limit CDK1 activity and halt mitotic progression. Subsequent degradation of CycB is required for anaphase progression and the production of two daughter cells in G1 phase of the cell cycle. During this transition from M phase back into G1 phase, RB is dephosphorylated, and the cycle is once more responsive to mitogenic and antiproliferative signalling. CCN, cyclin; CDC, cell division cycle; CDT1, chromatin licensing and DNA replication factor 1; MAD2L1, MAD2 mitotic arrest deficient-like 1; MCM, minichromosome maintenance complex component; PLK1, polo-like kinase 1.

The biology of CDKs

Integration of multiple signalling pathways through control of CDK4 and CDK6 activation. An understanding of the biology of CDKs is critical to deciphering the clinical results seen with CDK inhibitors, particularly in regard to determining biomarker and combination strategies. In most adult tissues, the majority of cells exit the cell cycle with diploid DNA content and are maintained in a quiescent G0 state. Tissue maintenance involves cues that physiologically induce cell cycle entry in a highly regulated manner. The mechanisms through which cells initiate entry into the cell cycle have been comprehensively described. Extracellular signals — including those activated by peptide growth factors (for example, RAS,

mitogen-activated protein kinase (MAPK) and mammalian target of rapamycin (mTOR)) and nuclear receptors (for example, the oestrogen receptor (ER) in mammary epithelia) — converge on the cell cycle to drive progression from G0 or G1 phase into S phase through regulation of the metazoan-specific CDK4 or CDK6 complex^{2,3,12,13}. CDK4 and CDK6 emerged phylogenetically with the appearance of multicellular organisms, and are subjected to multiple levels of regulation to control the transition into S phase. CDK4 and CDK6 are structurally related proteins that harbour many biochemical and biological similarities, although most published studies have focused on CDK4 (REF. 14). CDK6 is particularly important in promoting the proliferation of haematological precursors^{15,16}.

The activity of CDK4 and CDK6 is primarily controlled by their association with D-type cyclins (that is, cyclin D1, cyclin D2 and cyclin D3)^{17,18}. Among these, cyclin D1 is the best characterized. The expression of cyclin D1 is characterized as a 'delayed-early' response to mitogenic signalling, and intricate promoter and enhancer interactions control its transcription¹⁹. Although less well studied, cyclin D3 conforms to a similar pattern as cyclin D1, whereas the regulation of cyclin D2 remains more enigmatic, although cyclin D2 also drives proliferation in certain contexts^{20–24}. The differential expression of paralogues of D-type cyclins is likely to reflect tissue-specific aspects of normal physiology, wherein different D-type cyclins are expressed to promote CDK4 or CDK6 activation^{3,25}.

In addition to the transcriptional regulation of CDK4 and CDK6, the stability, intracellular localization and association of cyclin D with CDK4 and CDK6 are tightly regulated (FIG. 2a). In particular, cyclin D1 is unstable and actively shuttles between the cytoplasm and the nucleus. Phosphorylation of threonine 286 on cyclin D1 actively promotes its export and ubiquitin-mediated degradation^{26,27}. In contrast to other CDKs, for which cyclin association seems to occur relatively spontaneously, for CDK4 and CDK6 this process is regulated by multiple mechanisms²⁸. The inhibitor of CDK4 (INK4) proteins, which include p16^{INK4A}, p15^{INK4B}, p18^{INK4C} and p19^{INK4D}, represent CDK4- and CDK6-interacting proteins that seem to solely function as inhibitors of CDK4 and CDK6 (REFS 3,29,30). The INK4 proteins weaken the binding of D-type cyclins to CDK4 and CDK6, and also interact with the catalytic domains of CDK4 and CDK6 to potentially suppress kinase activity^{14,31,32}. These proteins therefore negatively regulate CDK4 and CDK6 in response to stress conditions³³. For example, p16^{INK4A} is induced by multiple oncoproteins to counteract transformation. Moreover, under stress conditions associated with cellular ageing³⁴, overexpression of p16^{INK4A} results in a profound G1 arrest of the cell cycle. Similarly, p15^{INK4B} is induced by transforming growth factor- β -mediated suppression of epithelial cell proliferation³⁵. CDK4 and CDK6, similar to other CDK proteins, are also subjected to phosphorylation^{36,37}. Thus, CDK4 and CDK6 serve as key nodes of integration downstream of multiple signalling pathways, in which their activation initiates progression into the cell cycle (FIG. 2a).

The association of D-type cyclins with CDK4 and CDK6 can induce kinase activity with a unique substrate spectrum compared with other CDKs³⁸. In particular, CDK4 and CDK6 have a specific preference for the phosphorylation of the tumour suppressor retinoblastoma protein (RB) and the related proteins p107 (also known as RBL1) and p130 (also known as RBL2)^{39,40} (FIG. 2b). RB, the first tumour suppressor identified, has been extensively studied^{41,42}. The RB protein does not have catalytic activity but functions through the assembly of multiprotein complexes to control the cell cycle. In particular, RB can bind to the E2F transcription factors, recruit co-repressors and repress the transcription of target genes that are regulated by E2Fs^{41,42}

(FIG. 2b). The E2Fs regulate the expression of a set of genes involved in cell cycle control (for example, cyclin E (*CCNE*), *CCNA* and *CCNB1*), dNTP biosynthesis (for example, dihydrofolate reductase (*DHFR*), ribonucleotide reductase M1 (*RRM1*) and *RRM2*) and mitotic progression (for example, polo-like kinase 1 (*PLK1*), BUB1 mitotic checkpoint serine/threonine kinase (*BUB1*) and spindle checkpoint protein MAD2 (*MAD2*)). The phosphorylation of RB by CDK4 and CDK6 initiates an intricate process of phosphorylation-mediated disruption of RB function that releases E2F and initiates subsequent progression through the cell cycle (FIG. 2c). CDK4 and CDK6 also phosphorylate forkhead box protein M1 (FOXO1), leading to stabilization of FOXO1 as a further mediator in the expression of genes required for progression through mitosis³⁸ (FIG. 3).

Deregulation of the CDK4/6–RB–p16^{INK4A} pathway in cancer. The CDK4/6–RB axis is critical to cell cycle entry; therefore, it is unsurprising that the vast majority of cancers subvert this axis to promote proliferation^{2,42,43} (FIG. 4). Most oncogenes promote the induction of p16^{INK4A} as an intrinsic break to deregulated proliferation^{34,44–46}. Overexpression of p16^{INK4A} ultimately engages RB to suppress growth and cell cycle progression, and promotes oncogene-induced senescence. Oncogene-induced senescence must be subverted to enable subsequent oncogenic proliferation, which occurs through two principal means in tumours: loss of p16^{INK4A} or loss of RB^{29,47}. Loss of p16^{INK4A} uncouples the oncogenic stress from the suppression of CDK4 or CDK6 activity, whereas loss of RB deregulates downstream signalling in the cell cycle. Consistent with this model, RB is required for the cell cycle arrest mediated by p16^{INK4A} (REFS 48,49). In addition, RB-negative tumours express super-physiological levels of p16^{INK4A} and are therefore insensitive to additional expression of p16^{INK4A} owing to the absence of RB²⁹.

A contrasting mechanism of deregulating the CDK4/6–RB axis is the direct oncogenic activation of CDK4 or CDK6 activity. Deregulated cyclin D1 protein expression, gene translocation and gene amplification are observed in many tumour types^{50–54}, and a plethora of functional data support the specific oncogenic activity of cyclin D1 (REFS 17,18,51). Furthermore, amplification of CDK4 and CDK6 is observed in several different types of cancer^{55,56}. Importantly, the distinct mechanisms of pathway dysregulation are mutually exclusive and are frequently tumour type-specific. For example, RB loss is a hallmark of small cell lung cancer, deregulation of cyclin D1 is common in breast cancer, and loss of p16^{INK4A} is particularly common in glioblastoma (FIG. 4).

Distal regulation of CDK2 and its deregulation in cancer. Although all CDKs have similarities, CDK2 is structurally and functionally related to CDK1 (REF. 3). CDK2 has a considerably broader substrate profile than CDK4 and CDK6, and it phosphorylates a large number of proteins involved in cell cycle progression (for example, p27^{KIP1} and RB), DNA replication (for example, replication factors A and C), histone synthesis (for example, NPAT), centrosome




Figure 2 | G1–S regulatory modules and relevance to cancer. Control over the G1–S transition is coordinated by distinct regulatory modules that are dysregulated in cancer. **a** | Initially, mitogenic signals impinge on cyclin-dependent kinase 4 (CDK4) or CDK6 activity through multiple mechanisms, including the induction of cyclin D1 (CycD1) gene (*CCND1*) expression, protein stability and assembly of the CDK–Cyc complex. These steps can be individually antagonized, or the induction of CDK4 and CDK6 inhibitors (that is, the inhibitor of CDK4 (INK4) family of proteins) can function to prevent complex assembly and to inhibit assembled complexes **b** | The net activation state of CDK4 and CDK6 initiates the phosphorylation of the tumour suppressor retinoblastoma protein (RB) that contributes to activation and release of the E2F family of transcription factors. E2F proteins control the expression of a multitude of positive-acting factors that are critical for progression through the S phase and the G2–M transition. Multiple mechanisms lead to RB inactivation in cancer, such as mutations, aberrant phosphorylation or protein sequestration. **c** | E2Fs and additional signals drive the expression and activation of CDK2–CycE and CDK2–CycA complexes, which contribute to DNA replication and further phosphorylation of RB. Deregulation of this activity is found in cancer through amplification of E-type cyclins or loss of CDK inhibitors. *CCN*, cyclin; *CDC*, cell division cycle; *CDT1*, chromatin licensing and DNA replication factor 1; *CIP*, CDK-interacting protein; *KIP*, kinase inhibitory protein; *MAD2L1*, MAD2 mitotic arrest deficient-like 1; *MCM*, minichromosome maintenance complex component; *PLK1*, polo-like kinase 1.

duplication (for example, nucleophosmin (NPM)), among other processes^{57–59} (FIG. 3). *In vitro*, CDK2 and its preferred E-type and A-type cyclin partners assemble spontaneously to form active kinase complexes^{3,60}. Much of the control over CDK2 involves the synthesis and availability of the cyclins, with RB and E2F regulating the abundance

of CDK2, cyclin E1 and cyclin E2 transcripts and proteins^{61–65}. This process couples mitogen-mediated activation of CDK4 and CDK6 with the activation of CDK2 (REFS 66,67) (FIG. 2c). In contrast to CDK4 and CDK6, CDK2 is not regulated by INK4 proteins^{30,68} but by the CDK-interacting protein/kinase inhibitory protein




Figure 3 | Summary of the biological functions of CDK complexes. A summary of the different classes of cyclin-dependent kinase (CDK)–cyclin (Cyc) complexes involved in the cell cycle or in diverse biological processes is shown. CDK–Cyc complexes shown in green promote cell cycle progression, whereas those depicted in blue are generally involved in transcriptional processes. The CDK5 complexes shown in red are involved both in the control of neuronal viability and in the promotion of the cell cycle. FOXM1, forkhead box protein M1; RB, retinoblastoma protein.

(CIP/KIP) class of CDK inhibitors, which bind to CDK2–cyclin complexes and render them inactive^{60,69–71}. For example, p21^{CIP1} acts as a DNA damage checkpoint (it is a critical downstream target gene of p53 that inhibits DNA synthesis), whereas p27^{KIP1} is responsive to mitogenic signalling as a further control on deregulated proliferation^{71,72}. Additionally, CDK2 is regulated by phosphorylation events⁷³. Therefore, multiple mechanisms in addition to the CDK4/6–RB axis can modulate the activity of CDK2 and subsequent DNA replication.

Recently, it has become clear that deregulation of CDK2 also occurs frequently in certain types of cancer⁷⁴. Cyclin E1 or cyclin E2 amplifications are key oncogenic events in multiple cancers, particularly in uterine and ovarian cancers^{75–77} (FIG. 4). Ectopic expression of cyclin E bypasses the need for CDK4 or CDK6 activity to initiate the S phase^{78–80}, and it is therefore assumed that amplification of E-type cyclins may be oncogenic in

a similar manner (that is, bypassing the physiological requirement for CDK4 or CDK6 activity to initiate expression of E-type cyclins). The CDK inhibitor p27^{KIP1} is downregulated in many cancers, although the genetic loss of p27^{KIP1} is fairly rare^{81,82}.

CDK1 is a key determinant of mitotic progression. CDK1 was cloned on the basis of complementarity to the *cdc2* gene of *Schizosaccharomyces pombe*⁸³. The expression of CDK1 and associated cyclins (cyclin A2 and cyclin B1) is coordinated through the activity of E2F-assembled complexes^{63,65}. These include the canonical E2F and RB constituents, as well as the transcription factor FOXM1 and the DREAM (dimerization partner, RB-like, E2F and multivulval class B) complex, which are particularly relevant for the coordination of transcripts involved in mitotic progression^{84–86}. The cyclins that assemble with CDK1 are intrinsically unstable and are regulated by ubiquitin-mediated degradation mechanisms, and the assembly and localization of CDK1 complexes are regulated by multiple overlapping mechanisms^{87–90}.

Mouse knockout experiments have shown that CDK1 is required for mammalian cell proliferation; it is the only CDK that can initiate the onset of mitosis (that is, the M phase)⁹¹. Premature initiation of mitosis before completion of the S phase results in chromosomal shattering and cell death⁹². Multiple factors restrain the activity of CDK1 until DNA replication is complete and there is minimal DNA damage. This integration of DNA replication and CDK1 activity is mediated by checkpoint signalling kinases such as CHK1 and WEE1, which suppress the activity of CDK1 via inhibitory phosphorylation⁹³, as well as through the cell division cycle 25 (CDC25) family of phosphatases. At the onset of mitosis, activation of CDK1 occurs rapidly through a positive feedback loop whereby CDK1 phosphorylates and inactivates WEE1. CDK1 subsequently phosphorylates multiple substrates, leading to nuclear envelope breakdown, chromosome condensation and mitotic spindle assembly⁹⁴ (FIG. 3). The subsequent progression from metaphase to anaphase is controlled by the spindle assembly checkpoints, and progression through anaphase is dependent on the attenuation of CDK1 activity through the degradation of cyclin B1 by the anaphase-promoting complex^{95,96}.

Interestingly, in contrast to the genetic deregulation of the CDKs that coordinate the S phase, there is limited evidence to show that CDK1 activity is dysregulated by direct genetic alteration in tumorigenesis. Derangement of p53 signalling or of DNA damage checkpoints indirectly leads to the deregulation of CDK1 (REFS 97,98), and high cyclin B1 expression is generally associated with a more aggressive cancer phenotype^{99,100}. However, the requirement that CDK1 activity must be attenuated to exit mitosis and the lethal aspects of uncoordinated CDK1 activity are likely to limit its potential as a direct oncogenic driver.

The role of cell cycle-independent CDKs. In addition to the well-known CDKs involved in regulating the cell cycle, there is an equivalent number of CDKs involved in

basal transcriptional regulation^{3,10,11} (FIG. 3). In particular, CDK8 is part of the mediator complex that regulates a plethora of genes^{101,102}. CDK7 has a general role in the phosphorylation of the RNA polymerase II carboxy-terminal domain that contributes to the initiation of transcription¹⁰³, and CDK9 also phosphorylates RNA polymerase II, thereby promoting elongation of transcription. Finally, CDK11 acts on the splicing machinery. In each of these contexts, the CDK activity is directed by specific cyclin interactions. Accumulating evidence suggests that these transcription-regulating CDKs may represent therapeutic targets in cancer.

In addition to the CDKs involved in transcriptional regulation, there remains a class of CDKs, including CDK3 and CDK5, for which the underlying functions are elusive. CDK3 was found to be intrinsically important for cell cycle control based on cell-based experiments that used a dominant-negative version of CDK3 (REF. 72) (FIG. 3). However, it was subsequently revealed that many mouse strains harbour an inactive CDK3, suggesting that its role in the cell cycle can be readily compensated¹⁰⁴. CDK5 was largely viewed as a neuronal kinase implicated in the protection of the nervous system from injury¹⁰⁵. However, recent work suggests that it harbours many functions similar to CDK4 and CDK2 in driving progression from G1–S and in RB phosphorylation in medullary thyroid cancer models¹⁰⁶. CDK5 might therefore have potential as a therapeutic target in this thyroid cancer subtype¹⁰⁶.

Development of pan-CDK inhibitors

The first generation of CDK inhibitors. Over the past 20 years, several CDK inhibitors have been developed as potential cancer therapeutics and tested in numerous trials and in several tumour types (FIG. 5). The first-generation CDK inhibitors developed were relatively nonspecific and may therefore be referred to as 'pan-CDK' inhibitors (an example of which is flavopiridol (also known as alvocidib; developed by Sanofi-Aventis), although some compounds, such as olomucine (not commercially developed) and roscovitine (also known as seliciclib; developed by Cyclacel), have relatively low affinity for CDK4 and CDK6). As the field of CDK biology progressed in parallel with the development of these agents, our understanding of their targets and interpretation of their behaviour have also evolved. For example, it was initially believed that roscovitine was a relatively specific inhibitor of CDK1, CDK2 and CDK5; however, subsequent data demonstrated that it also inhibited transcription, probably through the inhibition of CDK7 and CDK9 (REF. 107).

Of these first-generation inhibitors, flavopiridol is the most extensively investigated CDK inhibitor so far, with >60 clinical trials carried out between 1998 and 2014 (see [Supplementary information S1 \(box\)](#)). Flavopiridol is a semi-synthetic flavonoid derived from rohitukine, a chromone alkaloid, and has been shown to inhibit CDK1, CDK2, CDK4, CDK6, CDK7 and CDK9 (REFS 108,109). Although flavopiridol can induce cell cycle arrest in G1 and G2 phases, in certain contexts it also induces a cytotoxic response, probably as a result

of CDK7 and CDK9 inhibition that leads to suppression of transcription¹¹⁰. Although the underlying broad-spectrum nature of flavopiridol results in substantial *in vitro* activity, substantially less activity was observed *in vivo*¹¹⁰. Consequently, flavopiridol did not meet the initial high expectations for a CDK inhibitor, and low levels of clinical activity were seen in Phase II studies in several solid tumour types (see [Supplementary information S1 \(box\)](#)). There is evidence to indicate that flavopiridol may have clinical activity in haematological malignancies, such as chronic lymphocytic leukaemia (CLL) and mantle cell lymphoma^{111,112}, in which scheduling also seems to influence flavopiridol efficacy. For example, a relatively short infusion time (4 hours) resulted in response rates of 41% in 22 assessable patients with CLL^{113,114}. Patients with a high disease burden and high-risk genetic features achieved durable responses, and tumour lysis syndrome was reported in approximately 40% of patients with CLL treated with flavopiridol¹¹⁵. Despite these reports and extensive investment, no Phase III studies have emerged and drug development of flavopiridol was consequently discontinued in 2012.


In parallel with flavopiridol, roscovitine, a purine-based CDK inhibitor, was among the first agents to be evaluated in the clinic (see [Supplementary information S1 \(box\)](#)). Of the 56 patients treated in the Phase I setting, 1 patient achieved a partial response¹¹⁶. A subsequent blinded, randomized Phase II trial (APPRAISE) that compared roscovitine with the best supportive care was performed in patients with advanced non-small cell lung cancer. The APPRAISE study was terminated after 187 patients were enrolled; although results were never published, roscovitine did not seem to improve progression-free survival in this patient population (see [Cyclacel press release](#)). Currently, only a single trial is ongoing for roscovitine in Cushing disease.

Second-generation inhibitors of multiple CDKs.


Following on from flavopiridol and roscovitine, other CDK inhibitors were developed with the aim of increasing selectivity for CDK1 and CDK2 and/or increasing overall potency (FIG. 5). Again, numerous CDK inhibitors seemed to be particularly promising in preclinical studies, but only a few progressed past Phase I clinical trials^{117–120} (see [Supplementary information S1 \(box\)](#)).

Of these second-generation CDK inhibitors, dinaciclib (also known as MK-7965 and SCH727965; developed by Merck) has been most extensively studied in the clinic. Dinaciclib was specifically developed as a highly potent inhibitor of CDK1, CDK2, CDK5 and CDK9 (half-maximal inhibitory concentration (IC₅₀) values in the range of 1–4 nM), with less activity against CDK4, CDK6 and CDK7 (IC₅₀ values in the range of 60–100 nM). Compared to flavopiridol, dinaciclib exhibited superior activity with regard to suppression of RB phosphorylation in cell-based assays¹¹⁸. Moreover, dinaciclib inhibited cell cycle progression in >100 tumour cell lines of various tumour types and induced the regression of established solid tumours in a range of mouse models¹¹⁸. Initial results from Phase I studies were promising;


CDK4 and CDK6




CCND1





RB1



CDKN2A



CCNE1 and CCNE2



◀ **Figure 4 | Deregulation of CDK regulatory genes in cancer.** The frequencies of genetic amplification of cyclin-dependent kinase 4 (CDK4) and CDK6; cyclin D1 (CCND1); retinoblastoma 1 (RB1); cyclin-dependent kinase inhibitor 2A (CDKN2A); and cyclin E1 (CCNE1) and CCNE2 are summarized across multiple disease sites. For each of the indicated cancer types, the frequencies of mutation (green), amplification (red) and homozygous deletion (dark blue) were determined using genetic data from >2,000 cancer cases obtained through cBioPortal for Cancer Genomics. As shown, different types of cancer exhibit distinct predominant mechanisms of genetic alterations in cell cycle control. In many cases, the same cancer type has been evaluated in multiple independent studies. Detailed information about each case and disease is accessible through the cBioPortal for Cancer Genomics. ACC, adrenocortical carcinoma; ACYC, adenoid cystic carcinoma; AML, acute myeloid leukaemia; CCLE, Cancer Cell Line Encyclopedia; CS, carcinosarcoma; MBL, medulloblastoma; MM, multiple myeloma; NCI60, US National Cancer Institute (NCI) 60 human tumour cell line anticancer drug screen; SC, serous cystadenocarcinoma.

dinaciclib induced stable disease in a range of malignancies and displayed tolerable toxicity¹²¹. However, results from recent randomized Phase II trials of dinaciclib in solid tumours have been disappointing. A randomized Phase II trial of dinaciclib versus the chemotherapeutic agent capecitabine in advanced breast cancer was stopped after 30 patients were enrolled because interim analysis showed that the time to disease progression was inferior with dinaciclib treatment¹²². In a study evaluating dinaciclib monotherapy in patients with non-small cell lung cancer, dinaciclib showed no activity in previously treated patients¹²³. In adult patients with advanced acute myeloid leukaemia (patients ≥60 years of age) or acute lymphoblastic leukaemia, no objective responses were observed with dinaciclib¹²⁴. However, in patients with relapsed multiple myeloma, preliminary encouraging single-agent activity was reported in a Phase I/II trial, with 2 patients of 27 achieving partial responses¹²⁵, and dinaciclib, similar to flavopiridol, demonstrated clinical activity in pretreated patients with CLL¹²⁶. Based on these data, a randomized Phase III study of dinaciclib in refractory CLL is ongoing. Therefore, dinaciclib may prove useful in the treatment of certain haematological malignancies, in which flavopiridol also had evidence of activity.

Other CDK inhibitors include AT7519 (developed by Astex), a pyrazole 3-carboxamide compound that acts as an inhibitor of CDK1, CDK2, CDK4, CDK6 and CDK9. AT7519 has been evaluated in combination with bortezomib in a Phase II clinical trial enrolling patients with previously treated multiple myeloma. The combination was well tolerated, and more than one-third of patients achieved partial response¹²⁷. R547 (developed by Hoffman-La Roche) is an inhibitor of CDK1, CDK2 and CDK4 with less potency for CDK7, glycogen synthase kinase 3α (GSK3α) and GSK3β. R547 was tested in a Phase I study in 2007 as an intravenous weekly infusion¹²⁸. Although reported to have manageable side effects, there have not been further clinical trials with this compound. SNS-032 (also known as BMS-387032; developed by Bristol-Myers Squibb), which was initially described as having greater selectivity for CDK2 than CDK1 and CDK4, is now recognized to also target CDK7 and CDK9. Two Phase I clinical studies with

SNS-032, one in 2010 in advanced lymphoid malignancies¹²⁹ and one in 2008 in selected advanced tumours¹³⁰, have been reported, but no further developments are apparent. The development of AZD5438 (developed by Astra Zeneca) — an orally bioavailable, potent inhibitor of CDK1, CDK2 and CDK9 with less selectivity for CDK5 and CDK6 (REF. 131) — was discontinued after it was reported to be intolerable when administered continuously in patients with advanced solid tumours¹³². AG-024322 (developed by Pfizer) — a potent inhibitor of CDK1, CDK2 and CDK4 — was also discontinued in 2007 after the Phase I study was terminated, as it failed to achieve an acceptable clinical end point (see Supplementary information S1 (box)).

Reasons for failure of CDK inhibitors with low selectivity.

The general failure of non-selective CDK inhibitors in the clinic can be partly explained by at least three key underlying principles. First, there was a lack of clear understanding of the mechanism of action. For many of the CDK inhibitors with low specificity, there remains a lack of clarity with regard to which CDKs are actually being inhibited *in vivo* and therefore the corresponding mechanism that could underlie the therapeutic effect. For example, flavopiridol has been associated with diverse distal cellular effects, including cell cycle inhibition, transcriptional suppression, apoptosis, autophagy and endoplasmic reticulum stress^{133–135}. This lack of understanding confounds the ability to develop these agents as targeted therapies and to design effective combination strategies.

Second, there was a lack of appropriate patient selection. The vast majority of studies conducted with CDK inhibitors with low specificity were in unstratified patient cohorts. This is because there are essentially no biomarkers that may select for sensitive subpopulations for this class of inhibitors. The potential activity of both flavopiridol and dinaciclib in CLL and the rare ‘extraordinary’ responders suggest that there are molecular-based reasons that some tumours are vulnerable to such agents. Although the molecular underpinnings for these responses are unknown, it is tempting to speculate that the inhibition of CDKs that control transcription could underlie at least part of this activity.

Third, there is a lack of a therapeutic window. Many of these CDK inhibitors target several proteins that are critical to the proliferation (for example, CDK1) and survival (for example, CDK9) of normal cells. This limits the ability to achieve therapeutic levels of these drugs because of their intrinsic inability to discriminate between cancerous and healthy tissues. Consistent with this point, the toxicities associated with non-selective CDK inhibitors include diarrhoea, myelosuppression, anaemia and nausea^{116,121,136}.

A case for selectivity of CDK inhibitors. The ascribed weaknesses of pan-CDK inhibitors suggest that improved selectivity for certain CDKs is the key to the successful development of CDK inhibitors as therapeutic cancer agents. Selective inhibitors of CDK2 might provide the ability to target genetic and/or driver events

in tumours driven by cyclin E amplification. Emerging data suggest that targeting CDK1 is toxic in certain contexts, and it may be challenging to achieve a therapeutic window. For example, synthetic lethal screens against *KRAS* mutations have indicated a potential sensitivity

to CDK1 knockdown, although follow-up studies are required¹³⁷. Similarly, CDK1 or CDK9 inhibition is synthetically lethal with *MYC*^{138,139}. Pharmacologically, CDK1 inhibitors seem to potently cooperate with inhibitors of poly(ADP-ribose) polymerase¹⁴⁰ by




Figure 5 | **Selected CDK inhibitors.** The chemical structures of several pan-cyclin-dependent kinase (CDK) and CDK4- and CDK6-selective inhibitors are shown. The published half-maximal inhibitory concentration (IC₅₀) values against selected CDK complexes are shown. ND, not determined.

compromising DNA repair pathways. An alternative strategy is to selectively target CDK7, CDK8 or CDK9, which are associated with basal transcription, because cancer cells may harbour unique vulnerabilities to selective suppression. CDK8 may function as an oncoprotein in colorectal cancer by regulating the transcription of β -catenin target genes¹⁴¹. A covalent inhibitor of CDK7 (THZ1), which is relatively specific for CDK7 compared with other CDKs, has shown activity in multiple cancer cell models¹⁴². Similarly, a specific inhibitor for CDK9 (CDKI-73) that exhibited activity in animal models of leukaemia was recently developed¹⁴³.

Targeted inhibition of CDK4 and CDK6

Rationale for targeting CDK4 and CDK6. Based on the myriad findings from mechanistic studies and studies of CDK4 and CDK6 deregulation in cancer, three important features and expectations arose for CDK4 and CDK6 inhibitors in the clinic. First, one would expect that a pure CDK4 or CDK6 inhibitor would elicit a single phenotype in tumours: cytostatic G0/G1 arrest. Second, this effect would be a direct reflection of the engagement of RB to suppress gene expression and proliferation. Third, such effects would be particularly actionable in tumours that exhibit deregulation of CDK4 and CDK6 activity as opposed to other CDKs. Initial data from mouse models seeded confusion as to whether CDK4 and CDK6 were therapeutic targets because many tissues in the mouse developed normally despite the absence of CDK4 and/or CDK6 (REFS 15,144) and in the absence of D-type cyclins¹⁴⁵. These data reflected substantial compensatory plasticity with other CDKs. Despite uncertainties arising from the mouse knockout models, it subsequently became clear that attenuation of CDK4 and CDK6 activity could prevent the development of specific mouse tumour types. For example, cyclin D1 is crucial for the development of mammary tumours driven by HER2 (also known as ERBB2)¹⁴⁶, and similar observations were obtained in NOTCH-driven T cell leukaemia mouse models¹⁴⁷ and cell lines^{147,148}.

Pharmacological approaches. The development of selective inhibitors of both CDK4 and CDK6 has markedly changed the perception of CDKs as therapeutic targets in cancer. Through a combination of chemical screening and optimization, it was found that pyrido[2,3-*d*]pyrimidin-7-one compounds with a 2-aminopyridine side chain at the C2 position act as CDK inhibitors with a high degree of selectivity for CDK4 and CDK6 relative to other CDKs¹⁴⁹ (FIG. 5). Subsequent optimization resulted in the compound PD-0332991 (also known as palbociclib; developed by Pfizer) that induced potent G1 arrest in cell culture and xenograft models^{150,151}. As anticipated from the basic biology of G1–S transition, the effects of palbociclib were dependent on the presence of a functional RB protein, thus demonstrating a degree of biological specificity that had not been previously described for CDK inhibitors^{150,151}. Subsequently, multiple independent groups have demonstrated that specific CDK4

and CDK6 inhibitors arrest the cell cycle through the downstream blockade of phosphorylation of RB, as well as the related p107 and p130 proteins. This blockade results in the loss of expression of S-phase cyclins, nucleotide biosynthesis, DNA replication machinery and mitotic regulatory genes^{152,153}. Dual CDK4 and CDK6 inhibitors have been shown to be active in multiple preclinical models, including xenografts, genetically engineered mouse models and primary human tumour explants^{152,154–156} (TABLE 1). Parallel drug discovery efforts at Eli Lilly and Novartis resulted in the development of the drugs LY-2835219 (also known as abemaciclib) and LEE011, respectively^{157–162}. Each drug is structurally similar and chemically distinct from the less specific pan-CDK inhibitors (FIG. 5). The selectivity of all of these compounds is likely to reflect binding to the specialized ATP-binding pocket of CDK4 and CDK6 and specific interactions with residues in the ATP-binding cleft, although this has not been proven by structural analysis. In contrast to other CDK4- and CDK6-selective compounds, abemaciclib inhibits CDK9 in *in vitro* kinase assays, although there is no evidence of functional inhibition in cellular models¹⁵⁸.

Single-agent clinical outcomes. There are numerous emerging clinical studies with dual CDK4 and CDK6 inhibitors (TABLE 2). Results from a Phase I study with palbociclib monotherapy indicated promising clinical efficacy and a well-tolerated toxicity profile in patients with RB-positive advanced solid tumours and non-Hodgkin lymphoma. Of the 33 patients enrolled and treated with palbociclib (once daily for 14 days followed by 7 days off), 1 patient with testicular cancer achieved a partial response and 9 patients achieved stable disease¹⁶³. The anticipated cytostatic nature of dual CDK4 and CDK6 inhibitors resulted in prolonged stable disease duration of 18, 22 and 24 months in 3 men with growing teratoma syndrome with inoperable tumours¹⁶⁴. In patients with relapsed mantle cell lymphoma (which exhibits frequent cyclin D1 amplification) receiving palbociclib monotherapy, 5 of the 17 patients had >1 year progression-free survival, with 1 complete response and 2 partial responses¹⁶⁵. A Phase II study in liposarcoma, a disease with frequent CDK4 amplification, also reported favourable progression-free rates in patients with CDK4 amplification and RB expression^{165,166}. The recently reported Phase I study with LEE011 monotherapy in patients with advanced solid tumours demonstrated that LEE011 was well tolerated, with 2 confirmed partial responses and 40% of patients with stable disease¹⁶⁷. Similarly, abemaciclib exhibited single-agent activity associated with delayed disease progression and particularly robust activity in metastatic ER-positive breast cancer, although data are from a relatively small study with one group of patients¹⁶⁸. There are multiple ongoing Phase II trials evaluating dual CDK4 and CDK6 inhibitors as monotherapy in various tumour types (TABLE 2).

These early trials defined several key clinical hallmarks of inducing CDK4 and CDK6 inhibition in patients with cancer. Most importantly, it seems that neutropaenia is the principal dose-limiting toxicity of

Table 1 | **Preclinical outcome analysis of CDK4 and CDK6 inhibitors**

Indication	Agent	Cell culture	Animal model	Markers	Clinical	Refs
Mantle cell lymphoma	PD-0332991	Yes	Yes	–	Yes	165,187
Acute lymphoblastic lymphoma	PD-0332991	Yes	Yes	–	–	147,148
Multiple myeloma	PD-0332991	Yes	Yes	–	Yes	188
Liposarcoma	PD-0332991 and LEE011	Yes	Yes	CDK4	Yes	159,166
Hepatocellular carcinoma	PD-0332991	Yes	Yes	RB and p16 ^{INK4A}	Yes	153
Ovarian cancer	PD-0332991	Yes	Yes	RB and p16 ^{INK4A}	Yes	175
Breast cancer	PD-0332991, LEE-011	Yes	Yes	RB, p16 ^{INK4A} and luminal subtype	Yes	152,154, 155,169
Lung adenocarcinoma	PD-0332991	Yes	Yes	–	Yes	150,189
Prostate cancer	PD-0332991	Yes	Yes	–	Yes	190
Glioma	PD-0332991	Yes	Yes	RB and p16 ^{INK4A}	Yes	156,191,192
Renal cancer	PD-0332991	Yes	Yes	RB and p16 ^{INK4A}	–	176
Melanoma	PD-0332991 and LY2835219	Yes	Yes	–	Yes	183,193,194
Medulloblastoma	PD-0332991	Yes	Yes	–	–	195
Colon cancer	PD-0332991	Yes	Yes	–	Yes	150
Oesophageal cancer	PD-0332991	Yes	–	–	–	196
Neuroblastoma	LEE011	Yes	Yes	–	Yes	157
Pancreatic cancer	PD-0332991	Yes	Yes	–	–	184,197

CDK, cyclin-dependent kinase; RB, retinoblastoma protein.

palbociclib and LEE011. Although neutropaenia is a common side effect of cytotoxic agents, the neutropaenia associated with palbociclib and LEE011 is distinct in that it is rapidly reversible, reflecting a cytostatic effect on neutrophil precursors in the bone marrow. Consequently, both palbociclib and LEE011 are dosed intermittently to accommodate a break for haematological recovery¹⁶⁷. Interestingly, abemaciclib exhibits more prominent gastrointestinal-associated toxicity, whereas neutropaenia is less evident, enabling continuous dosing. The reasons behind these observations and implications for future development remain unclear.

Hormone therapy combination strategies. Preclinical investigation suggested that dual CDK4 and CDK6 inhibitors positively interact with other therapeutic agents. In particular, synergy was observed when palbociclib was combined with hormone therapy in ER-positive breast cancer cell lines, although the observed effects can range from additive to synergistic depending on the model¹⁵⁵. Additionally, CDK4 and CDK6 inhibition has shown activity in multiple ER-positive breast cancer models that have acquired resistance to ER antagonists^{155,169,170}. Importantly, resistance to endocrine therapy is associated with the

deregulation of proliferation-associated genes that are regulated by the CDK4/6–RB–E2F axis, suggesting a basis for cooperativity in the clinic^{43,171}. These observations triggered a series of randomized Phase II studies that have consequently transformed the CDK field.

The PALOMA-1 Phase II clinical trial randomized 165 women with advanced ER-positive breast cancer into two treatment groups: the aromatase inhibitor letrozole versus letrozole plus palbociclib. Data from this trial showed that the combination of palbociclib plus letrozole elicited significant improvement in median progression-free survival compared with letrozole alone (20.2 months compared with 10.2 months; hazard ratio (HR) = 0.488 (95% CI: 0.319–0.748) and one-sided $p = 0.0004$). The overall survival analysis after 61 deaths demonstrated a trend in favour of the letrozole plus palbociclib combination (37.5 months versus 33.3 months, respectively; HR = 0.813; $p = 0.2105$)^{172–174}. Consequently, palbociclib received Breakthrough Therapy designation by the US Food and Drug Administration in April 2013. ER-positive breast cancer is characterized by frequent dysregulation of CDK4 and CDK6 activity due to the overexpression and amplification of the gene encoding cyclin D1 (*CCND1*). Those cancers with amplification of *CCND1* seemed to derive no greater benefit from

Table 2 | Reported clinical trials with targeted CDK4 and CDK6 inhibitors

Tumour type	Phase	Dosage	Response rate	Refs
Palbociclib (PD-0332991)				
Advanced melanoma, breast cancer, renal cancer, ovarian cancer, liposarcoma and colon cancer, among others	<ul style="list-style-type: none"> Phase Ia (dose escalation) N=41 	<ul style="list-style-type: none"> Administered over 6 cohorts (standard 3+3) 3 weeks on, 1 week off RP2D: 125 mg PO OD 	SD: 27% (10/37)	178
Liposarcoma, colon cancer and melanoma, among others	<ul style="list-style-type: none"> Phase Ia N=33 (RB positive) 	<ul style="list-style-type: none"> Administered over 4 cohorts 2 weeks on; 1 week off RP2D: 200 mg PO OD 	<ul style="list-style-type: none"> PR: 3% (1/31; testicular cancer) SD: 29% (9/31) 	163
Relapsed mantle cell lymphoma with ≥1 of: CCND1 positivity by immunostaining, t(11;14) translocation on cytogenetic analysis and molecular evidence of CCND1-IGH rearrangement	<ul style="list-style-type: none"> Phase I N=17 	<ul style="list-style-type: none"> 100–125 mg PO OD 	<ul style="list-style-type: none"> CR: 6% (1/16) PR: 12% (2/16) SD: 43% (7/16) 	165
ER-positive and HER2-negative metastatic breast cancer	<ul style="list-style-type: none"> Phase Ib N=12 	<ul style="list-style-type: none"> Palbociclib (125 mg PO OD; 2 weeks on; 1 week off) + letrozole (2.5 mg PO OD; continuous) 	<ul style="list-style-type: none"> PR: 25% (3/12) SD: 75% (9/12) 	198
ER-positive and HER2-negative metastatic breast cancer	<ul style="list-style-type: none"> Phase II (PALOMA-1; TRIO-18) Palbociclib + letrozole versus letrozole alone in 1:1 randomization N=165 	<ul style="list-style-type: none"> Palbociclib (125 mg; 3 weeks on; 1 week off) + letrozole (2.5 mg; continuous) 	<ul style="list-style-type: none"> PFS: 20.2 months for palbociclib + letrozole versus 10.2 months for letrozole alone (HR=0.488 (95% CI: 0.319–0.748) and 1-sided p=0.0004) OS: 37.5 months for palbociclib + letrozole versus 33.3 months for letrozole alone (HR=0.813; p=0.2105) 	173
Metastatic breast cancer (64% ER-positive; 7% ER-positive and HER2-positive; 29% TNBC)	<ul style="list-style-type: none"> Phase II N=37 (RB positive) 	<ul style="list-style-type: none"> 125 mg PO 3 weeks on; 1 week off 	<ul style="list-style-type: none"> PR: 7% (2/28) SD: 50% (14/28) PFS: 4.1 months for ER-positive (95% CI: 2.3–7.7) PFS: 18.8 months for ER-positive and HER2-positive (95% CI: 5.1–∞) PFS: 1.8 months (95% CI 0.9–∞) for TNBC 	199
Well-differentiated (17%) and dedifferentiated (83%) liposarcoma with CDK4 amplification detected by FISH and RB expression detected by IHC	<ul style="list-style-type: none"> Phase II N=30 	<ul style="list-style-type: none"> 200 mg PO OD 2 weeks on; 1 week off 	<ul style="list-style-type: none"> PR: 3% (1/30) at 74 weeks; 19/30 were progression-free at 12 weeks PFS: median 18 weeks PFS: 66% (90% CI: 51–100%) at 12 weeks Met primary end point of exceeding PFS rate of 40% at 12 weeks for active second-line agent 	166
LEE011				
RB-positive advanced solid tumours and lymphomas	<ul style="list-style-type: none"> Phase I N=132 	<ul style="list-style-type: none"> Stage 1 (N=85): Treatment arm 1: escalating LEE011 doses (3 weeks on; 1 week off) Treatment arm 2: escalating LEE011 doses (continuous) Stage 2 (N=47): RP2D expansion MTD: 900 mg; RP2D: 600 mg using 3 weeks on; 1 week off schedule 	<ul style="list-style-type: none"> PR: 2.9% (2/70) at 600 mg per day SD: 26% with >4 cycles and 14% with >6 cycles 	167
Post-menopausal ER-positive, HER2-negative metastatic breast cancer	<ul style="list-style-type: none"> Phase Ib LEE011 + everolimus (mTOR inhibitor) + exemestane (aromatase inhibitor) N=6 	<ul style="list-style-type: none"> Treatment arm 1: escalating LEE011 doses (starting 200 mg per day; 3 weeks on; 1 week off) + everolimus (2.5 mg per day, fixed continuous) + exemestane (25 mg per day; continuous) Treatment arm 2: safety run-in with LEE011 (600 mg per day; 3 weeks on; 1 week off) + exemestane (25 mg per day; continuous) 	<ul style="list-style-type: none"> Preliminary results indicate that triple combination is tolerable Efficacy data not yet available 	200

Table 2 (cont.) | Reported clinical trials with targeted CDK4 and CDK6 inhibitors

Tumour type	Phase	Dosage	Response rate	Refs
Post-menopausal ER-positive, HER2-negative locally advanced or metastatic breast cancer	<ul style="list-style-type: none"> Phase Ib LEE011 + BYL719 (PI3Kα inhibitor) + letrozole N = 11 	<ul style="list-style-type: none"> Treatment arm 1: LEE011 (3 weeks on; 1 week off) + letrozole (2.5 mg; continuous); 4 week cycle Treatment arm 2: BYL719 (continuous) + letrozole (2.5 mg; continuous); 4 week cycle Treatment arm 3: LEE011 + BYL719 (continuous) + letrozole (2.5 mg; continuous); 4 week cycle 	Efficacy data not yet available	201
NRAS-mutant metastatic melanoma	<ul style="list-style-type: none"> Phase Ib (single arm) LEE011 + binimetinib (MEK inhibitor) N = 14 	<ul style="list-style-type: none"> LEE011 (starting 200 mg per day OD; 3 weeks on; 1 week off) + binimetinib (45 mg PO BD) 	<ul style="list-style-type: none"> PR: 43% (6/14) SD: 43% (6/14) Promising preliminary antitumour activity 	202
Abemaciclib (LY2835219)				
Non-small cell lung cancer (KRAS wild type and KRAS mutant)	<ul style="list-style-type: none"> Phase I N = 49 	<ul style="list-style-type: none"> MTD already established at 200 mg in earlier stage of study Treatment arm 1 (N = 25): 200 mg PO BD continuous (28-day cycle) Treatment arm 2 (N = 24): 150 mg PO BD continuous (28-day cycle) 	<ul style="list-style-type: none"> RR: 2% PR (1/49) Overall DCR: 51% DCR 37% (19/49) for KRAS wild type and 54% (26/49) for KRAS mutant PFS: 2.1 months 	203
Hormone receptor-positive metastatic breast cancer	<ul style="list-style-type: none"> Phase I Abemaciclib + fulvestrant N = 60 	<ul style="list-style-type: none"> Treatment arm 1 (N = 47): abemaciclib (200 mg BD PO; continuous; 28-day cycle) Treatment arm 2 (N = 13): abemaciclib + fulvestrant (500 mg IM every 4 weeks) 	<ul style="list-style-type: none"> PR: 17% (8/47) with 6% (3/47) unconfirmed Single-agent activity demonstrated; acceptable safety profile in combination with fulvestrant Further evaluation required 	204

BD, twice daily; CCND1, cyclin D1; CDK, cyclin-dependent kinase; CR, complete response; DCR, disease control rate (CR+PR+SD); ER, oestrogen receptor; FISH, fluorescence *in situ* hybridization; HR, hazard ratio; IHC, immunohistochemistry; IGH, immunoglobulin heavy locus; IM, intramuscularly; MTD, maximum tolerated dose; mTOR, mammalian target of rapamycin; N, number of patients; OD, once daily; OS, overall survival; PFS, progression-free survival; PI3K α , phosphatidylinositol-4,5-bisphosphate 3-kinase (PI3K), catalytic subunit- α ; PO, oral route; PR, partial response; RB, retinoblastoma protein; RP2D, recommended Phase II dose; RR, response rate; SD, stable disease; TNBC, triple-negative breast cancer.

palbociclib, an observation that is likely to reflect the central nature of cyclin D1 in promoting ER-positive breast cancer proliferation regardless of whether high *CCND1* expression was due to amplification or other mechanisms.

As a follow-up to these findings, multiple Phase II and III trials of combination therapies were initiated. The combination therapies tested each include a dual CDK4 and CDK6 inhibitor (abemaciclib, LEE011 or palbociclib) and a hormone therapy (letrozole, anastrozole or fulvestrant) (TABLE 2).

CDK4 and CDK6 inhibitor biomarker strategies.

Preclinical work has defined a series of biomarkers that may be used in the selection of tumours that may respond to dual CDK4 and CDK6 inhibitors. The most conservative and best supported of these markers is the direct assessment of the CDK4–RB–p16^{INK4A} pathway. Data from multiple groups have demonstrated that RB is necessary for the arrest induced by CDK4 and CDK6 inhibition^{152–157,175,176}, and loss of RB is therefore a marker of resistance to CDK4 and CDK6 inhibition. Loss of RB results in supra-physiological expression of p16^{INK4A}, which may also be a biomarker of resistance. For example, high levels of p16^{INK4A} are identified in malignancies caused by human papilloma virus, a virus that can inactivate RB in cervical and in head and neck cancers^{29,177}.

Whether dual assessment of RB loss and induction of p16^{INK4A} expression is better than either biomarker alone is uncertain. There remains a considerable need to identify other predictive markers for tumours with CDK4 and/or CDK6 dependence or ‘addiction’ that can be selectively targeted. Examples of other predictive markers could be the amplification of cyclin D1 or CDK4 and CDK6, loss of p16^{INK4A} or other genetic alterations leading to the deregulation of CDK4 or CDK6 activity. This concept has been incorporated into ‘basket trial’ designs with palbociclib (LUNG-MAP) and LEE011 (SIGNATURE), in which patients with specific signature mutations that would be expected to deregulate CDK4 and CDK6 activity can be enrolled for treatment with these inhibitors.

The future of CDK4 and CDK6 inhibitors. Preclinical and clinical data suggest that dual CDK4 and CDK6 inhibitors could have broad-ranging efficacy in many cancer indications. Several questions have arisen from the published work regarding understanding which diseases would benefit the most from dual CDK4 and CDK6 inhibitors.

One important question is how to determine whether an RB-proficient tumour benefits from CDK4 and CDK6 inhibition. In some tumour types, CDK4 and CDK6 inhibition has a surprisingly modest clinical effect despite

molecular alterations indicating a robust response^{163,178}. Some cancer types seem either to be innately resistant or to acquire rapid resistance to the effects of CDK4 and CDK6 inhibition. For example, CDK4 and CDK6 suppression seems to have little clinical effect in colorectal cancer, triple-negative breast cancer and melanomas. Therefore, such tumours would not benefit from monotherapy in the absence of potent combination strategies and robust predictive markers. In these cancers, other CDKs, particularly CDK2, are likely to compensate for selective CDK4 and CDK6 inhibition. However, the factors that determine whether other CDKs can compensate for CDK4 and CDK6 inhibition are poorly understood.

Another question is how to optimize schedules for treatment. CDK4 and CDK6 inhibition antagonizes the effect of cytotoxic chemotherapy and radiotherapy because the vast majority of cytotoxic chemotherapies require cells to be cycling^{179–181}. Despite this requirement, several studies are underway to evaluate scheduling with CDK4 and CDK6 inhibition, following the concept that release from CDK4 and CDK6 inhibition may synchronize cells and thereby sensitize cancer cells to a subsequent cytotoxic treatment, or may prevent ongoing proliferation or re-population of cancer cells between cytotoxic administrations¹⁸². Although such scheduling approaches have been shown to be potentially beneficial in preclinical models, translating this to the clinic, where proliferation rates of tumours are highly variable, will be challenging. A variation of this principle in patients with a known RB-inactivated cancer is the potential use of CDK4 and CDK6 inhibition to protect normal cells from the effect of chemotherapy or radiotherapy while rendering the tumour vulnerable¹⁷⁹.

Finally, it is important to determine ideal combinations. Considerable interest lies in the potential for combining CDK4 and CDK6 inhibitors with other targeted

therapies. Substantial preclinical work has demonstrated that CDK4 and CDK6 inhibition may be synergistic with MEK inhibition in *NRAS*-positive melanoma¹⁸³. Similarly, in pancreatic cancer, CDK4 and CDK6 inhibition is synergistic with inhibitors of insulin-like growth factor 1 receptor and mTOR^{184,185}. Furthermore, in cancers with mutated phosphatidylinositol-4,5-bisphosphate 3-kinase (PI3K), catalytic subunit- α (PIK3CA), PI3K inhibitors may synergize with CDK4 and CDK6 inhibitors¹⁶⁰. Studies with LEE011 have incorporated this CDK4 and CDK6 inhibitor into the triplet combinations with a PIK3CA inhibitor (BYL719) and letrozole ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT01872260) identifier: NCT01872260), and the mTOR inhibitor everolimus with exemestane in ER-positive breast cancers (NCT01857193). Recently released data from a study in melanoma suggest that such rational combinations are effective and that CDK4 and CDK6 inhibition could represent a preferred combination agent with a range of targeted agents¹⁸⁶.

Conclusion

CDK complexes have critical roles in multiple aspects of biology, including proliferation control and transcription. After the generally disappointing results seen in clinical trials with non-selective CDK inhibitors, the importance of selectivity of compounds for specific CDKs and of patient selection is now widely accepted. The main challenges will be in the development of a suite of highly selective agents against specific CDKs, companion diagnostics that will enable the selection of appropriate patient populations, and a firm understanding of the intersection of pharmacology and biology that will provide the basis for rational drug combinations. Now, >10 years after Hunt, Nurse and Hartwell were awarded the Nobel Prize for the identification of CDKs, the promise of their seminal studies is finally beginning to be realized.

- Nurse, P., Masui, Y. & Hartwell, L. Understanding the cell cycle. *Nature Med.* **4**, 1103–1106 (1998).
- Sherr, C. J. Cancer cell cycles. *Science* **274**, 1672–1677 (1996).
- Lim, S. & Kaldis, P. Cdks, cyclins and CKIs: roles beyond cell cycle regulation. *Development* **140**, 3079–3093 (2013).
- Malumbres, M. Therapeutic opportunities to control tumor cell cycles. *Clin. Transl. Oncol.* **8**, 399–408 (2006).
- Hunt, T., Nasmyth, K. & Novak, B. The cell cycle. *Phil. Trans. R. Soc. B* **366**, 3494–3497 (2011).
- Hunt, T. Nobel Lecture. Protein synthesis, proteolysis, and cell cycle transitions. *Biosci. Rep.* **22**, 465–486 (2002).
- Nurse, P. M. Nobel Lecture. Cyclin dependent kinases and cell cycle control. *Biosci. Rep.* **22**, 487–499 (2002).
- Hartwell, L. H. Nobel Lecture. Yeast and cancer. *Biosci. Rep.* **22**, 373–394 (2002).
References 6–8 provide highly insightful overviews of the Nobel Prize-winning findings that underpin much of the subsequent analyses of cell cycle regulatory processes.
- Drapkin, R., Le Roy, G., Cho, H., Akoulitchev, S. & Reinberg, D. Human cyclin-dependent kinase-activating kinase exists in three distinct complexes. *Proc. Natl Acad. Sci. USA* **93**, 6488–6493 (1996).
- Bregman, D. B., Pestell, R. G. & Kidd, V. J. Cell cycle regulation and RNA polymerase II. *Front. Biosci.* **5**, D244–D257 (2000).
- Nemet, J., Jellic, B., Rubelj, I. & Sopta, M. The two faces of Cdk8, a positive/negative regulator of transcription. *Biochimie* **97**, 22–27 (2014).
- Malumbres, M. & Barbacid, M. To cycle or not to cycle: a critical decision in cancer. *Nature Rev. Cancer* **1**, 222–231 (2001).
- Rodgers, J. T. *et al.* mTORC1 controls the adaptive transition of quiescent stem cells from G0 to G(Alert). *Nature* **510**, 393–396 (2014).
- Pavletich, N. P. Mechanisms of cyclin-dependent kinase regulation: structures of CDKs, their cyclin activators, and CIP and INK4 inhibitors. *J. Mol. Biol.* **287**, 821–828 (1999).
- Malumbres, M. *et al.* Mammalian cells cycle without the D-type cyclin-dependent kinases Cdk4 and Cdk6. *Cell* **118**, 493–504 (2004).
This is one of multiple papers demonstrating that selective CDK activities can be bypassed by compensatory pathways and the underlying plasticity within the cell cycle.
- Hu, M. G. *et al.* A requirement for cyclin-dependent kinase 6 in thymocyte development and tumorigenesis. *Cancer Res.* **69**, 810–818 (2009).
- Sherr, C. J. D-type cyclins. *Trends Biochem. Sci.* **20**, 187–190 (1995).
- Diehl, J. A. Cycling to cancer with cyclin D1. *Cancer Biol. Ther.* **1**, 226–231 (2002).
- Matsushime, H., Roussel, M. F., Ashmun, R. A. & Sherr, C. J. Colony-stimulating factor 1 regulates novel cyclins during the G1 phase of the cell cycle. *Cell* **65**, 701–713 (1991).
- Spofford, L. S., Abel, E. V., Boisvert-Adamo, K. & Aplin, A. E. Cyclin D3 expression in melanoma cells is regulated by adhesion-dependent phosphatidylinositol 3-kinase signaling and contributes to G1–S progression. *J. Biol. Chem.* **281**, 25644–25651 (2006).
- Sicinski, P. *et al.* Cyclin D2 is an FSH-responsive gene involved in gonadal cell proliferation and oncogenesis. *Nature* **384**, 470–474 (1996).
- Kushner, J. A. *et al.* Cyclins D2 and D1 are essential for postnatal pancreatic β -cell growth. *Mol. Cell. Biol.* **25**, 3752–3762 (2005).
- Yu, Q., Ciernerych, M. A. & Sicinski, P. Ras and Myc can drive oncogenic cell proliferation through individual D-cyclins. *Oncogene* **24**, 7114–7119 (2005).
- Cooper, A. B. *et al.* A unique function for cyclin D3 in early B cell development. *Nature Immunol.* **7**, 489–497 (2006).
- Ciernerych, M. A. *et al.* Development of mice expressing a single D-type cyclin. *Genes Dev.* **16**, 3277–3289 (2002).
- Diehl, J. A., Zindy, F. & Sherr, C. J. Inhibition of cyclin D1 phosphorylation on threonine-286 prevents its rapid degradation via the ubiquitin-proteasome pathway. *Genes Dev.* **11**, 957–972 (1997).

27. Diehl, J. A., Cheng, M., Roussel, M. F. & Sherr, C. J. Glycogen synthase kinase-3 β regulates cyclin D1 proteolysis and subcellular localization. *Genes Dev.* **12**, 3499–3511 (1998).
28. Cheng, M., Sexl, V., Sherr, C. J. & Roussel, M. F. Assembly of cyclin D-dependent kinase and titration of p27^{Kip1} regulated by mitogen-activated protein kinase kinase (MEK1). *Proc. Natl Acad. Sci. USA* **95**, 1091–1096 (1998).
29. Witkiewicz, A. K., Knudsen, K. E., Dicker, A. P. & Knudsen, E. S. The meaning of p16^{INK4a} expression in tumors: functional significance, clinical associations and future developments. *Cell Cycle* **10**, 2497–2503 (2011).
30. Serrano, M., Hannon, G. J. & Beach, D. A new regulatory motif in cell-cycle control causing specific inhibition of cyclin D/CDK4. *Nature* **366**, 704–707 (1993).
31. Jeffrey, P. D., Tong, L. & Pavletich, N. P. Structural basis of inhibition of CDK–cyclin complexes by INK4 inhibitors. *Genes Dev.* **14**, 3115–3125 (2000).
32. Russo, A. A., Tong, L., Lee, J. O., Jeffrey, P. D. & Pavletich, N. P. Structural basis for inhibition of the cyclin-dependent kinase Cdk6 by the tumour suppressor p16^{INK4a}. *Nature* **395**, 237–243 (1998).
33. Serrano, M. & Blasco, M. A. Putting the stress on senescence. *Curr. Opin. Cell Biol.* **13**, 748–753 (2001).
34. Serrano, M., Lin, A. W., McCurrach, M. E., Beach, D. & Lowe, S. W. Oncogenic ras provokes premature cell senescence associated with accumulation of p53 and p16^{INK4a}. *Cell* **88**, 593–602 (1997).
- This study provides a key link between the induction of p16^{INK4a} and the blockade of oncogene-driven tumorigenesis.**
35. Reynisdottir, I., Polyak, K., Iavarone, A. & Massague, J. Kip/Cip and INK4 CDK inhibitors cooperate to induce cell cycle arrest in response to TGF- β . *Genes Dev.* **9**, 1831–1845 (1995).
36. Terada, Y., Tatsuka, M., Jinno, S. & Okayama, H. Requirement for tyrosine phosphorylation of Cdk4 in G1 arrest induced by ultraviolet irradiation. *Nature* **376**, 358–362 (1995).
37. Bertero, T. et al. CDC25A targeting by miR-483-3p decreases CCND-CDK4/6 assembly and contributes to cell cycle arrest. *Cell Death Differ.* **20**, 800–811 (2013).
38. Anders, L. et al. A systematic screen for CDK4/6 substrates links FOXM1 phosphorylation to senescence suppression in cancer cells. *Cancer Cell* **20**, 620–634 (2011).
39. Kato, J., Matsushime, H., Hiebert, S. W., Ewen, M. E. & Sherr, C. J. Direct binding of cyclin D to the retinoblastoma gene product (pRb) and pRb phosphorylation by the cyclin D-dependent kinase CDK4. *Genes Dev.* **7**, 331–342 (1993).
40. Matsushime, H. et al. D-type cyclin-dependent kinase activity in mammalian cells. *Mol. Cell. Biol.* **14**, 2066–2076 (1994).
41. Wang, J. Y., Knudsen, E. S. & Welch, P. J. The retinoblastoma tumor suppressor protein. *Adv. Cancer Res.* **64**, 25–85 (1994).
42. Burkhardt, D. L. & Sage, J. Cellular mechanisms of tumour suppression by the retinoblastoma gene. *Nature Rev. Cancer* **8**, 671–682 (2008).
43. Knudsen, E. S. & Knudsen, K. E. Tailoring to RB: tumour suppressor status and therapeutic response. *Nature Rev. Cancer* **8**, 714–724 (2008).
44. Bartkova, J. et al. Oncogene-induced senescence is part of the tumorigenesis barrier imposed by DNA damage checkpoints. *Nature* **444**, 633–637 (2006).
45. Michaloglou, C. et al. BRAFE600-associated senescence-like cell cycle arrest of human naevi. *Nature* **436**, 720–724 (2005).
46. Burd, C. E. et al. Monitoring tumorigenesis and senescence *in vivo* with a p16^{INK4a}-luciferase model. *Cell* **152**, 340–351 (2013).
47. LaPak, K. M. & Burd, C. E. The molecular balancing act of p16^{INK4a} in cancer and aging. *Mol. Cancer Res.* **12**, 167–183 (2014).
48. Lukas, J. et al. Retinoblastoma-protein-dependent cell-cycle inhibition by the tumour suppressor p16. *Nature* **375**, 503–506 (1995).
49. Lukas, J., Bartkova, J., Rohde, M., Strauss, M. & Bartek, J. Cyclin D1 is dispensable for G1 control in retinoblastoma gene-deficient cells independently of cdk4 activity. *Mol. Cell. Biol.* **15**, 2600–2611 (1995).
50. Motokura, T. et al. A novel cyclin encoded by a bcl1-linked candidate oncogene. *Nature* **350**, 512–515 (1991).
51. Knudsen, K. E., Diehl, J. A., Haiman, C. A. & Knudsen, E. S. Cyclin D1: polymorphism, aberrant splicing and cancer risk. *Oncogene* **25**, 1620–1628 (2006).
52. Jiang, W. et al. Amplification and expression of the human cyclin D gene in esophageal cancer. *Cancer Res.* **52**, 2980–2983 (1992).
53. Buckley, M. F. et al. Expression and amplification of cyclin genes in human breast cancer. *Oncogene* **8**, 2127–2133 (1993).
54. Bartkova, J. et al. Cyclin D1 protein expression and function in human breast cancer. *Int. J. Cancer* **57**, 353–361 (1994).
55. Khatib, Z. A. et al. Coamplification of the CDK4 gene with MDM2 and GLI in human sarcomas. *Cancer Res.* **53**, 5535–5541 (1993).
56. Park, S. et al. Aberrant CDK4 amplification in refractory rhabdomyosarcoma as identified by genomic profiling. *Sci. Rep.* **4**, 3623 (2014).
57. Ma, T. et al. Cell cycle-regulated phosphorylation of p220(NPAT) by cyclin E/Cdk2 in Cajal bodies promotes histone gene transcription. *Genes Dev.* **14**, 2298–2313 (2000).
58. Okuda, M. et al. Nucleophosmin/B23 is a target of CDK2/cyclin E in centrosome duplication. *Cell* **103**, 127–140 (2000).
59. Sever-Chroneos, Z. et al. Retinoblastoma tumor suppressor protein signals through inhibition of cyclin-dependent kinase 2 activity to disrupt PCNA function in S phase. *Mol. Cell. Biol.* **21**, 4032–4045 (2001).
60. Sherr, C. J. & Roberts, J. M. CDK inhibitors: positive and negative regulators of G1-phase progression. *Genes Dev.* **13**, 1501–1512 (1999).
61. Herrera, R. E. et al. Altered cell cycle kinetics, gene expression, and G1 restriction point regulation in Rb-deficient fibroblasts. *Mol. Cell. Biol.* **16**, 2402–2407 (1996).
62. Zhu, W., Giangrande, P. H. & Nevins, J. R. E2Fs link the control of G1/S and G2/M transcription. *EMBO J.* **23**, 4615–4626 (2004).
63. Markey, M. P. et al. Unbiased analysis of RB-mediated transcriptional repression identifies novel targets and distinctions from E2F action. *Cancer Res.* **62**, 6587–6597 (2002).
64. Moroy, T. & Geisen, C. Cyclin, E. *Int. J. Biochem. Cell Biol.* **36**, 1424–1439 (2004).
65. Ren, B. et al. E2F integrates cell cycle progression with DNA repair, replication, and G₂/M checkpoints. *Genes Dev.* **16**, 245–256 (2002).
66. Sherr, C. J. Cell cycle control and cancer. *Harvey Lect.* **96**, 73–92 (2000).
67. Bartek, J., Bartkova, J. & Lukas, J. The retinoblastoma protein pathway in cell cycle control and cancer. *Exp. Cell Res.* **237**, 1–6 (1997).
68. Xiong, Y., Zhang, H. & Beach, D. Subunit rearrangement of the cyclin-dependent kinases is associated with cellular transformation. *Genes Dev.* **7**, 1572–1583 (1993).
69. el-Deiry, W. S. et al. WAF1, a potential mediator of p53 tumor suppression. *Cell* **75**, 817–825 (1993).
70. Polyak, K. et al. p27Kip1, a cyclin-Cdk inhibitor, links transforming growth factor- β and contact inhibition to cell cycle arrest. *Genes Dev.* **8**, 9–22 (1994).
71. Polyak, K. et al. Cloning of p27Kip1, a cyclin-dependent kinase inhibitor and a potential mediator of extracellular antimetagenic signals. *Cell* **78**, 59–66 (1994).
72. van den Heuvel, S. & Harlow, E. Distinct roles for cyclin-dependent kinases in cell cycle control. *Science* **262**, 2050–2054 (1993).
73. Wu, C. L. et al. Cables enhances cdk2 tyrosine 15 phosphorylation by Wee1, inhibits cell growth, and is lost in many human colon and squamous cancers. *Cancer Res.* **61**, 7325–7332 (2001).
74. Scaltriti, M. et al. Cyclin E amplification/overexpression is a mechanism of trastuzumab resistance in HER2⁺ breast cancer patients. *Proc. Natl Acad. Sci. USA* **108**, 3761–3766 (2011).
75. Etemadmoghadam, D. et al. Synthetic lethality between CCNE1 amplification and loss of BRCA1. *Proc. Natl Acad. Sci. USA* **110**, 19489–19494 (2013).
76. Karst, A. M. et al. Cyclin E1 deregulation occurs early in secretory cell transformation to promote formation of fallopian tube-derived high-grade serous ovarian cancers. *Cancer Res.* **74**, 1141–1152 (2014).
77. Kuhn, E., Bahadiri-Talbot, A. & Shih, I. E. M. Frequent CCNE1 amplification in endometrial intraepithelial carcinoma and uterine serous carcinoma. *Mod. Pathol.* **27**, 1014–1019 (2014).
78. Caldon, C. E. et al. Cyclin E2 overexpression is associated with endocrine resistance but not insensitivity to CDK2 inhibition in human breast cancer cells. *Mol. Cancer Ther.* **11**, 1488–1499 (2012).
79. Lukas, J. et al. Cyclin E-induced S phase without activation of the pRB/E2F pathway. *Genes Dev.* **11**, 1479–1492 (1997).
80. Knudsen, E. S., Buckmaster, C., Chen, T. T., Feramisco, J. R. & Wang, J. Y. Inhibition of DNA synthesis by RB: effects on G1/S transition and S-phase progression. *Genes Dev.* **12**, 2278–2292 (1998).
81. Hershko, D. D. Cyclin-dependent kinase inhibitor p27 as a prognostic biomarker and potential cancer therapeutic target. *Future Oncol.* **6**, 1837–1847 (2010).
82. Chu, I. M., Hengst, L. & Slingerland, J. M. The CDK inhibitor p27 in human cancer: prognostic potential and relevance to anticancer therapy. *Nature Rev. Cancer* **8**, 253–267 (2008).
83. Draetta, G., Brizuela, L., Potashkin, J. & Beach, D. Identification of p34 and p13, human homologs of the cell cycle regulators of fission yeast encoded by cdc2⁺ and suc1⁺. *Cell* **50**, 319–325 (1987).
84. Sadasivam, S., Duan, S. & DeCaprio, J. A. The MuvB complex sequentially recruits B-Myb and FoxM1 to promote mitotic gene expression. *Genes Dev.* **26**, 474–489 (2012).
85. Sadasivam, S. & DeCaprio, J. A. The DREAM complex: master coordinator of cell cycle-dependent gene expression. *Nature Rev. Cancer* **13**, 585–595 (2013).
86. Wang, I. C. et al. Forkhead box M1 regulates the transcriptional network of genes essential for mitotic progression and genes encoding the SCF (Skp2–Cks1) ubiquitin ligase. *Mol. Cell. Biol.* **25**, 10875–10894 (2005).
87. Draviam, V. M., Orrechia, S., Lowe, M., Pardi, R. & Pines, J. The localization of human cyclins B1 and B2 determines CDK1 substrate specificity and neither enzyme requires MEK to disassemble the Golgi apparatus. *J. Cell Biol.* **152**, 945–958 (2001).
88. Gavet, O. & Pines, J. Progressive activation of CyclinB1–Cdk1 coordinates entry to mitosis. *Dev. Cell* **18**, 533–543 (2010).
89. Gavet, O. & Pines, J. Activation of cyclin B1–Cdk1 synchronizes events in the nucleus and the cytoplasm at mitosis. *J. Cell Biol.* **189**, 247–259 (2010).
90. Nigg, E. A. Polo-like kinases: positive regulators of cell division from start to finish. *Curr. Opin. Cell Biol.* **10**, 776–783 (1998).
91. Santamaria, D. et al. Cdk1 is sufficient to drive the mammalian cell cycle. *Nature* **448**, 811–815 (2007).
92. Aarts, M. et al. Forced mitotic entry of S-phase cells as a therapeutic strategy induced by inhibition of WEE1. *Cancer Discov.* **2**, 524–539 (2012).
93. Nigg, E. A. Mitotic kinases as regulators of cell division and its checkpoints. *Nature Rev. Mol. Cell Biol.* **2**, 21–32 (2001).
94. Nigg, E. A., Blangy, A. & Lane, H. A. Dynamic changes in nuclear architecture during mitosis: on the role of protein phosphorylation in spindle assembly and chromosome segregation. *Exp. Cell Res.* **229**, 174–180 (1996).
95. Bhargadwaj, R. & Yu, H. The spindle checkpoint, aneuploidy, and cancer. *Oncogene* **23**, 2016–2027 (2004).
96. Morgan, D. O. Regulation of the APC and the exit from mitosis. *Nature Cell Biol.* **1**, E47–E53 (1999).
97. Taylor, W. R. & Stark, G. R. Regulation of the G2/M transition by p53. *Oncogene* **20**, 1803–1815 (2001).
98. Bayart, E., Grigorieva, O., Leibovitch, S., Onclercq-Delic, R. & Amor-Gueret, M. A major role for mitotic CDC2 kinase inactivation in the establishment of the mitotic DNA damage checkpoint. *Cancer Res.* **64**, 8954–8959 (2004).
99. Aaltonen, K. et al. High cyclin B1 expression is associated with poor survival in breast cancer. *Br. J. Cancer* **100**, 1055–1060 (2009).
100. Nimeus-Malmstrom, E. et al. Cyclin B1 is a prognostic proliferation marker with a high reproducibility in a population-based lymph node negative breast cancer cohort. *Int. J. Cancer* **127**, 961–967 (2010).

101. Tassan, J. P., Jaquenoud, M., Leopold, P., Schultz, S. J. & Nigg, E. A. Identification of human cyclin-dependent kinase 8, a putative protein kinase partner for cyclin C. *Proc. Natl Acad. Sci. USA* **92**, 8871–8875 (1995).
102. Akoulitchev, S., Chuiikov, S. & Reinberg, D. TFIH is negatively regulated by CDK8-containing mediator complexes. *Nature* **407**, 102–106 (2000). **This seminal study defines the role of selected CDK family members in transcriptional regulation.**
103. Spangler, L., Wang, X., Conaway, J. W., Conaway, R. C. & Dvir, A. TFIH action in transcription initiation and promoter escape requires distinct regions of downstream promoter DNA. *Proc. Natl Acad. Sci. USA* **98**, 5544–5549 (2001).
104. Ye, X., Zhu, C. & Harper, J. W. A premature-termination mutation in the *Mus musculus* cyclin-dependent kinase 3 gene. *Proc. Natl Acad. Sci. USA* **98**, 1682–1686 (2001).
105. Ohshima, T. et al. Targeted disruption of the cyclin-dependent kinase 5 gene results in abnormal corticogenesis, neuronal pathology and perinatal death. *Proc. Natl Acad. Sci. USA* **93**, 11173–11178 (1996).
106. Pozo, K. et al. The role of CDK5 in neuroendocrine thyroid cancer. *Cancer Cell* **24**, 499–511 (2013). **This important study defines a role for CDK5 in driving tumour initiation and progression and as a potential therapeutic target in cancer.**
107. Whittaker, S. R., Walton, M. I., Garrett, M. D. & Workman, P. The Cyclin-dependent kinase inhibitor CYC202 (R-roscovitine) inhibits retinoblastoma protein phosphorylation, causes loss of Cyclin D1, and activates the mitogen-activated protein kinase pathway. *Cancer Res.* **64**, 262–272 (2004).
108. Shapiro, G. I. Cyclin-dependent kinase pathways as targets for cancer treatment. *J. Clin. Oncol.* **24**, 1770–1783 (2006).
109. Sedlacek, H. et al. Flavopiridol (L86 8275; NSC 649890), a new kinase inhibitor for tumor therapy. *Int. J. Oncol.* **9**, 1143–1168 (1996).
110. Bose, P., S. G. L. & Grant, S. Cyclin-dependent kinase inhibitor therapy for hematologic malignancies. *Expert Opin. Investig. Drugs* **22**, 723–738 (2013).
111. Lin, T. S. et al. Flavopiridol, fludarabine, and rituximab in mantle cell lymphoma and indolent B-cell lymphoproliferative disorders. *J. Clin. Oncol.* **28**, 418–423 (2010).
112. Kouroukis, C. T. et al. Flavopiridol in untreated or relapsed mantle-cell lymphoma: results of a phase II study of the National Cancer Institute of Canada Clinical Trials Group. *J. Clin. Oncol.* **21**, 1740–1745 (2003).
113. Byrd, J. C. et al. Flavopiridol administered using a pharmacologically derived schedule is associated with marked clinical efficacy in refractory, genetically high-risk chronic lymphocytic leukemia. *Blood* **109**, 399–404 (2007).
114. Lapenna, S., Giordano, A. Cell cycle kinases as therapeutic targets for cancer. *Nature Rev. Drug Discov.* **8**, 547–566 (2009).
115. Blum, K. A. et al. Risk factors for tumor lysis syndrome in patients with chronic lymphocytic leukemia treated with the cyclin-dependent kinase inhibitor, flavopiridol. *Leukemia* **25**, 1444–1451 (2011).
116. Le Tourneau, C. et al. Phase I evaluation of seliciclib (R-roscovitine), a novel oral cyclin-dependent kinase inhibitor, in patients with advanced malignancies. *Eur. J. Cancer* **46**, 3243–3250 (2010).
117. Payton, M. et al. Discovery and evaluation of dual CDK1 and CDK2 inhibitors. *Cancer Res.* **66**, 4299–4308 (2006).
118. Parry, D. et al. Dinaciclib (SCH 727965), a novel and potent cyclin-dependent kinase inhibitor. *Mol. Cancer Ther.* **9**, 2344–2353 (2010). **This study illustrates the potency of the multiple-CDK inhibitor dinaciclib in preclinical models.**
119. DePinto, W. et al. *In vitro* and *in vivo* activity of R547: a potent and selective cyclin-dependent kinase inhibitor currently in Phase I clinical trials. *Mol. Cancer Ther.* **5**, 2644–2658 (2006).
120. Misra, R. N. et al. N-(cycloalkylamino) acyl-2-aminothiazole inhibitors of cyclin-dependent kinase 2. N-[5-[[[5-(1,1-dimethylethyl)-2-oxazolyl] methyl]thio]-2-thiazolyl]-4- piperidinecarboxamide (BMS-387032), a highly efficacious and selective antitumor agent. *J. Med. Chem.* **47**, 1719–1728 (2004).
121. Nemunaitis, J. J. et al. A first-in-human, Phase 1, dose-escalation study of dinaciclib, a novel cyclin-dependent kinase inhibitor, administered weekly in subjects with advanced malignancies. *J. Transl. Med.* **11**, 259 (2013).
122. Mita, M. M. et al. Randomized Phase II trial of the cyclin-dependent kinase inhibitor dinaciclib (MK-7965) versus capecitabine in patients with advanced breast cancer. *Clin. Breast Cancer* **14**, 169–176 (2014).
123. Stephenson, J. J. et al. Randomized Phase 2 study of the cyclin-dependent kinase inhibitor dinaciclib (MK-7965) versus erlotinib in patients with non-small cell lung cancer. *Lung Cancer* **83**, 219–223 (2014).
124. Gojo, I. et al. Phase II study of the cyclin-dependent kinase (CDK) inhibitor dinaciclib (SCH 727965) in patients with advanced acute leukemias. *ASH Annual Meeting Abstracts* **116**, 3287 (2010).
125. Kumar, S. K. et al. Phase 1/2 trial of a novel CDK inhibitor dinaciclib (SCH727965) in patients with relapsed multiple myeloma demonstrates encouraging single agent activity. *ASH Annual Meeting Abstracts* **120**, 76 (2012).
126. Blachly, J. S. & Byrd, J. C. Emerging drug profile: cyclin-dependent kinase inhibitors. *Leuk. Lymphoma* **54**, 2135–2143 (2013).
127. Raju, N. et al. A Phase I/II open-label multicenter study of the cyclin kinase inhibitor AT7519M alone and in combination with bortezomib in patients with previously treated multiple myeloma (ASH 55th meeting abstract). *Blood* **122**, 1976 (2013).
128. Diab, S. et al. A Phase I study of R547, a novel, selective inhibitor of cell cycle and transcriptional cyclin dependent kinases (CDKs). *J. Clin. Oncol.* **25**, 3528 (2007).
129. Tong, W. G. et al. Phase I and pharmacologic study of SNS-032, a potent and selective Cdk 2, 7, and 9 inhibitor, in patients with advanced chronic lymphocytic leukemia and multiple myeloma. *J. Clin. Oncol.* **28**, 3015–3022 (2010).
130. Heath, E. L., B. K., Martell, R. E., Adelman, D. C. & Lorusso, P. M. A phase I study of SNS-032 (formerly BMS-387032), a potent inhibitor of cyclin-dependent kinases 2, 7 and 9 administered as a single oral dose and weekly infusion in patients with metastatic refractory solid tumors. *Invest. New Drugs* **26**, 59–65 (2008).
131. Byth, K. F. et al. AZD5438, a potent oral inhibitor of cyclin-dependent kinases 1, 2, and 9, leads to pharmacodynamic changes and potent antitumor effects in human tumor xenografts. *Mol. Cancer Ther.* **8**, 1856–1866 (2009).
132. Boss, D. S. et al. Safety, tolerability, pharmacokinetics and pharmacodynamics of the oral cyclin-dependent kinase inhibitor AZD5438 when administered at intermittent and continuous dosing schedules in patients with advanced solid tumours. *Ann. Oncol.* **21**, 884–894 (2010).
133. Mahoney, E., Byrd, J. C. & Johnson, A. J. Autophagy and ER stress play an essential role in the mechanism of action and drug resistance of the cyclin-dependent kinase inhibitor flavopiridol. *Autophagy* **9**, 434–435 (2013).
134. Carlson, B. A., Dubay, M. M., Sausville, E. A., Brizuela, L. & Worland, P. J. Flavopiridol induces G1 arrest with inhibition of cyclin-dependent kinase (CDK) 2 and CDK4 in human breast carcinoma cells. *Cancer Res.* **56**, 2973–2978 (1996).
135. Kelland, L. R. Flavopiridol, the first cyclin-dependent kinase inhibitor to enter the clinic: current status. *Expert Opin. Investig. Drugs* **9**, 2903–2911 (2000).
136. Senderowicz, A. M. et al. Phase I trial of continuous infusion flavopiridol, a novel cyclin-dependent kinase inhibitor, in patients with refractory neoplasms. *J. Clin. Oncol.* **16**, 2986–2999 (1998).
137. Barbie, D. A. et al. Systematic RNA interference reveals that oncogenic KRAS-driven cancers require TBK1. *Nature* **462**, 108–112 (2009).
138. Kang, J., Sergio, C. M., Sutherland, R. L. & Musgrove, E. A. Targeting cyclin-dependent kinase 1 (CDK1) but not CDK4/6 or CDK2 is selectively lethal to MYC-dependent human breast cancer cells. *BMC Cancer* **14**, 32 (2014).
139. Huang, C. H. et al. CDK9-mediated transcription elongation is required for MYC addiction in hepatocellular carcinoma. *Genes Dev.* **28**, 1800–1814 (2014).
140. Johnson, N. et al. Compromised CDK1 activity sensitizes BRCA-proficient cancers to PARP inhibition. *Nature Med.* **17**, 875–882 (2011).
141. Firestein, R. et al. CDK8 is a colorectal cancer oncogene that regulates β -catenin activity. *Nature* **455**, 547–551 (2008).
142. Kwiatkowski, N. et al. Targeting transcription regulation in cancer with a covalent CDK7 inhibitor. *Nature* **511**, 616–620 (2014).
143. Walsby, E. et al. A novel Cdk9 inhibitor preferentially targets tumor cells and synergizes with fludarabine. *Oncotarget* **5**, 375–385 (2014).
144. Kozar, K. & Sicinski, P. Cell cycle progression without cyclin D–CDK4 and cyclin D–CDK6 complexes. *Cell Cycle* **4**, 388–391 (2005).
145. Kozar, K. et al. Mouse development and cell proliferation in the absence of D-cyclins. *Cell* **118**, 477–491 (2004).
146. Yu, Q., Geng, Y. & Sicinski, P. Specific protection against breast cancers by cyclin D1 ablation. *Nature* **411**, 1017–1021 (2001). **This seminal study illustrates the tumour context-specific requirement for cyclin D1.**
147. Choi, Y. J. et al. The requirement for cyclin D function in tumor maintenance. *Cancer Cell* **22**, 438–451 (2012).
148. Sawai, C. M. et al. Therapeutic targeting of the cyclin D3:CDK4/6 complex in T cell leukemia. *Cancer Cell* **22**, 452–465 (2012).
149. VanderWel, S. N. et al. Pyrido[2,3-d]pyrimidin-7-ones as specific inhibitors of cyclin-dependent kinase 4. *J. Med. Chem.* **48**, 2371–2387 (2005).
150. Fry, D. W. et al. Specific inhibition of cyclin-dependent kinase 4/6 by PD 0332991 and associated antitumor activity in human tumor xenografts. *Mol. Cancer Ther.* **3**, 1427–1438 (2004).
151. Toogood, P. L. et al. Discovery of a potent and selective inhibitor of cyclin-dependent kinase 4/6. *J. Med. Chem.* **48**, 2388–2406 (2005).
152. Dean, J. L., Thangavel, C., McClelland, A. K., Reed, C. A. & Knudsen, E. S. Therapeutic CDK4/6 inhibition in breast cancer: key mechanisms of response and failure. *Oncogene* **29**, 4018–4032 (2010).
153. Rivadeneira, D. B. et al. Proliferative suppression by CDK4/6 inhibition: complex function of the retinoblastoma pathway in liver tissue and hepatoma cells. *Gastroenterology* **138**, 1920–1930 (2010).
154. Dean, J. L. et al. Therapeutic response to CDK4/6 inhibition in breast cancer defined by *ex vivo* analyses of human tumors. *Cell Cycle* **11**, 2756–2761 (2012).
155. Finn, R. S. et al. PD 0332991, a selective cyclin D kinase 4/6 inhibitor, preferentially inhibits proliferation of luminal estrogen receptor-positive human breast cancer cell lines *in vitro*. *Breast Cancer Res.* **11**, R77 (2009).
156. Michaud, K. et al. Pharmacologic inhibition of cyclin-dependent kinases 4 and 6 arrests the growth of glioblastoma multiforme intracranial xenografts. *Cancer Res.* **70**, 3228–3238 (2010).
157. Rader, J. et al. Dual CDK4/CDK6 inhibition induces cell-cycle arrest and senescence in neuroblastoma. *Clin. Cancer Res.* **19**, 6173–6182 (2013).
158. Dickson, M. A. Molecular pathways: CDK4 inhibitors for cancer therapy. *Clin. Cancer Res.* **20**, 3379–3383 (2014).
159. Zhang, Y. X. et al. Antiproliferative effects of CDK4/6 inhibition in CDK4-amplified human liposarcoma *in vitro* and *in vivo*. *Mol. Cancer Ther.* **13**, 2184–2193 (2014).
160. Vora, S. R. et al. CDK 4/6 inhibitors sensitize PIK3CA mutant breast cancer to PI3K inhibitors. *Cancer Cell* **26**, 136–149 (2014).
161. Gelbert, L. M. et al. Preclinical characterization of the CDK4/6 inhibitor LY2835219: *in-vivo* cell cycle-dependent/independent anti-tumor activities alone/in combination with gemcitabine. *Invest. New Drugs* **32**, 825–837 (2014).
162. Tate, S. C. et al. Semi-mechanistic pharmacokinetic/Pharmacodynamic modeling of the antitumor activity of LY2835219, a new cyclin-dependent kinase 4/6 inhibitor, in mice bearing human tumor xenografts. *Clin. Cancer Res.* **20**, 3763–3774 (2014).
163. Schwartz, G. K. et al. Phase I study of PD 0332991, a cyclin-dependent kinase inhibitor, administered in 3-week cycles (Schedule 2/1). *Br. J. Cancer* **104**, 1862–1868 (2011).
164. Vaughn, D. et al. Treatment of growing teratoma syndrome. *N. Engl. J. Med.* **360**, 423–424 (2009).
165. Leonard, J. P. et al. Selective CDK4/6 inhibition with tumor responses by PD0332991 in patients with mantle cell lymphoma. *Blood* **119**, 4597–4607 (2012).

This Phase II study provides proof of concept that selective CDK4 and CDK6 inhibitors can have clinical activity in selected tumours.

166. Dickson, M. A. *et al.* Phase II trial of the CDK4 inhibitor PD0332991 in patients with advanced CDK4-amplified well-differentiated or dedifferentiated liposarcoma. *J. Clin. Oncol.* **31**, 2024–2028 (2013).
167. Infante, J. R. *et al.* A Phase I study of the single-agent CDK4/6 inhibitor LEE011 in pts with advanced solid tumors and lymphomas. *J. Clin. Oncol.* **32** (Suppl.), 2528 (2014).
168. Shapiro, G. *et al.* A first-in-human phase I study of the CDK4/6 inhibitor, LY2835219, for patients with advanced cancer. *J. Clin. Oncol.* **31** (Suppl.), 2500, (2013).
169. Miller, T. W. *et al.* ER α -dependent E2F transcription can mediate resistance to estrogen deprivation in human breast cancer. *Cancer Discov.* **1**, 338–351 (2011).
170. Thangavel, C. *et al.* Therapeutically activating RB: reestablishing cell cycle control in endocrine therapy-resistant breast cancer. *Endocr. Relat. Cancer* **18**, 333–345 (2011).
- References 155, 169 and 170 provide much of the evidence that CDK4 and CDK6 inhibitors would be efficacious in ER-positive breast cancer.**
171. Desmedt, C. & Sotiriou, C. Proliferation: the most prominent predictor of clinical outcome in breast cancer. *Cell Cycle* **5**, 2198–2202 (2006).
172. Finn, R. S. *et al.* S-16 Results of a randomized phase 2 study of PD 0332991, a cyclin-dependent kinase (CDK) 4/6 inhibitor, in combination with letrozole vs letrozole alone for first-line treatment of ER $^{+}$ /HER2 $^{-}$ advanced breast cancer (BC). *Cancer Res.* **72** (Suppl. 24) (2012).
173. Finn, R. S. *et al.* Final results of a randomized Phase II study of PD 0332991, a cyclin-dependent kinase (CDK) 4/6 inhibitor, in combination with letrozole versus letrozole alone for first-line treatment of ER $^{+}$ /HER2 $^{-}$ advanced breast cancer (PALOMA-1; TRIO-18). *Cancer Res.* **74**, CT101 (2014).
174. Finn, R. S. *et al.* The cyclin-dependent kinase 4/6 inhibitor palbociclib in combination with letrozole versus letrozole alone as first-line treatment of oestrogen receptor-positive, HER2-negative, advanced breast cancer (PALOMA-1/TRIO-18): a randomised phase 2 study. *Lancet Oncol.* **16**, 25–35 (2015).
- This paper presents the clinical data that now support multiple Phase III studies of CDK4 and CDK6 inhibition in concert with hormone therapy.**
175. Konecny, G. E. *et al.* Expression of p16 and retinoblastoma determines response to CDK4/6 inhibition in ovarian cancer. *Clin. Cancer Res.* **17**, 1591–1602 (2011).
176. Logan, J. E. *et al.* PD-0332991, a potent and selective inhibitor of cyclin-dependent kinase 4/6, demonstrates inhibition of proliferation in renal cell carcinoma at nanomolar concentrations and molecular markers predict for sensitivity. *Anticancer Res.* **33**, 2997–3004 (2013).
177. Carozzi, F. *et al.* Use of p16-INK4A overexpression to increase the specificity of human papillomavirus testing: a nested substudy of the NTCC randomised controlled trial. *Lancet Oncol.* **9**, 937–945 (2008).
178. Flaherty, K. T. *et al.* Phase I, dose-escalation trial of the oral cyclin-dependent kinase 4/6 inhibitor PD 0332991, administered using a 21-day schedule

in patients with advanced cancer. *Clin. Cancer Res.* **18**, 568–576 (2012).

179. Roberts, P. J. *et al.* Multiple roles of cyclin-dependent kinase 4/6 inhibitors in cancer therapy. *J. Natl Cancer Inst.* **104**, 476–487 (2012).
180. McClendon, A. K. *et al.* CDK4/6 inhibition antagonizes the cytotoxic response to anthracycline therapy. *Cell Cycle* **11**, 2747–2755 (2012).
181. Dean, J. L., McClendon, A. K. & Knudsen, E. S. Modification of the DNA damage response by therapeutic CDK4/6 inhibition. *J. Biol. Chem.* **287**, 29075–29087 (2012).
182. Witkiewicz, A. K., Cox, D. & Knudsen, E. S. CDK4/6 inhibition provides a potent adjunct to Her2-targeted therapies in preclinical breast cancer models. *Genes Cancer* **5**, 261–272 (2014).
183. Kwong, L. N. *et al.* Oncogenic NRAS signaling differentially regulates survival and proliferation in melanoma. *Nature Med.* **18**, 1503–1510 (2012).
184. Heilmann, A. M. *et al.* CDK4/6 and IGF1 receptor inhibitors synergize to suppress the growth of p16^{INK4A}-deficient pancreatic cancers. *Cancer Res.* **74**, 3947–3958 (2014).
185. Franco, J., Witkiewicz, A. K. & Knudsen, E. S. CDK4/6 inhibitors have potent activity in combination with pathway selective therapeutic agents in models of pancreatic cancer. *Oncotarget* **5**, 6512–6525 (2014).
186. [No authors listed.] Drug combo shows promise in NRAS-mutant melanoma. *Cancer Discov.* <http://dx.doi.org/10.1158/2159-8290.CD-NB2014-098> (2014).
- References 160 and 181–185 demonstrate the important cooperative mechanisms between CDK4 and CDK6 and targeted therapies in multiple cancers.**
187. Marzec, M. *et al.* Mantle cell lymphoma cells express predominantly cyclin D1a isoform and are highly sensitive to selective inhibition of CDK4 kinase activity. *Blood* **108**, 1744–1750 (2006).
188. Menu, E. *et al.* A novel therapeutic combination using PD 0332991 and bortezomib: study in the ST33MM myeloma model. *Cancer Res.* **68**, 5519–5523 (2008).
189. Puyol, M. *et al.* A synthetic lethal interaction between K-RAS oncogenes and CDK4 unveils a therapeutic strategy for non-small cell lung carcinoma. *Cancer Cell* **18**, 63–73 (2010).
190. Comstock, C. E. *et al.* Targeting cell cycle and hormone receptor pathways in cancer. *Oncogene* **32**, 5481–5491 (2013).
191. Barton, K. L. *et al.* PD-0332991, a CDK4/6 inhibitor, significantly prolongs survival in a genetically engineered mouse model of brainstem glioma. *PLoS ONE* **8**, e77639 (2013).
192. Wiedemeyer, W. R. *et al.* Pattern of retinoblastoma pathway inactivation dictates response to CDK4/6 inhibition in GBM. *Proc. Natl Acad. Sci. USA* **107**, 11501–11506 (2010).
193. Young, R. J. *et al.* Loss of CDKN2A expression is a frequent event in primary invasive melanoma and correlates with sensitivity to the CDK4/6 inhibitor PD0332991 in melanoma cell lines. *Pigment Cell. Melanoma Res.* **27**, 590–600 (2014).
194. Yadav, V. *et al.* The CDK4/6 inhibitor LY2835219 overcomes vemurafenib resistance resulting from MAPK reactivation and cyclin D1 upregulation. *Mol. Cancer Ther.* **13**, 2253–2263 (2014).
195. Whiteway, S. L. *et al.* Inhibition of cyclin-dependent kinase 6 suppresses cell proliferation and enhances

radiation sensitivity in medulloblastoma cells. *J. Neurooncol* **111**, 113–121 (2013).

196. Ismail, A. *et al.* Early G₁ cyclin-dependent kinases as prognostic markers and potential therapeutic targets in esophageal adenocarcinoma. *Clin. Cancer Res.* **17**, 4513–4522 (2011).
197. Liu, F. & Korc, M. Cdk4/6 inhibition induces epithelial-mesenchymal transition and enhances invasiveness in pancreatic cancer cells. *Mol. Cancer Ther.* **11**, 2138–2148 (2012).
198. Slamon, D. J. *et al.* Phase I study of PD 0332991, cyclin-D kinase (CDK) 4/6 inhibitor in combination with letrozole for first-line treatment of patients with ER-positive, HER2-negative breast cancer. *J. Clin. Oncol. Abstr.* **28**:15s, 3060 (2010).
199. DeMichele, A. *et al.* A Phase II trial of an oral CDK 4/6 inhibitor, D0332991, in advanced breast cancer. *J. Clin. Oncol. Abstr.* **31**, 519 (2013).
200. Bardia, A. *et al.* Phase Ib/II study of LEE011, everolimus, and exemestane in postmenopausal women with ER $^{+}$ /HER2-metastatic breast cancer. *J. Clin. Oncol.* **32** (Suppl.), 535 (2014).
201. Munster, P. *et al.* Phase Ib study of LEE011 and BYL719 in combination with letrozole in estrogen receptor-positive, HER2-negative breast cancer (ER $^{+}$, HER2 $^{-}$ BC). *J. Clin. Oncol.* **32** (Suppl.), 533 (2014).
202. Sosman, J. *et al.* A phase 1b/2 study of LEE011 in combination with binimetinib (MEK162) in patients with NRAS-mutant melanoma: early encouraging clinical activity. *J. Clin. Oncol.* **32** (Suppl.), 9009 (2014).
203. Goldman, J.W. *et al.* Clinical activity of LY2835219, a novel cell cycle inhibitor selective for CDK4 and CDK6, in patients with non-small cell lung cancer. *J. Clin. Oncol.* **32** (Suppl.), 8026 (2014).
204. Patnaik, A. *et al.* LY2825219, a novel cell cycle inhibitor for CDK4/6, in combination with fulvestrant for patients with hormone receptor positive (HR + ve) metastatic breast cancer. *J. Clin. Oncol.* **32** (Suppl.), 524 (2014).

Acknowledgements

The authors thank their colleagues in the field for guidance, critical discussion and editorial advice related to the preparation of this manuscript. They also regret any omissions. N.C.T. acknowledges funding from the UK National Health Service to the Royal Marsden and the Institute of Cancer Research NIHR Biomedical Research Centre. E.S.K. acknowledges research funding from the US National Institutes of Health/National Cancer Institute (NIH/NCI) (CA129134 and CA188650). A.K.W. acknowledges research funding from the NIH/NCI (CA163863).

Competing interests statement

The authors declare **competing interests**: see Web version for details.

FURTHER INFORMATION

cBioPortal for Cancer Genomics: <http://www.cbioportal.org/public-portal>

ClinicalTrials.gov: <https://clinicaltrials.gov>

Cyclacel press release: <http://investor.cyclacel.com/releasedetail.cfm?ReleaseID=614395>

SUPPLEMENTARY INFORMATION

See online article: **S1** (box)

ALL LINKS ARE ACTIVE IN THE ONLINE PDF